

A Contribution to the Phenomenological Theory of Kinetic Processes in Ferromagnetic Dielectrics. I. The Relaxation in the Gas of the Spin Waves SOV/56-34-6-30/51

Akhiyezer, L. D. Landau and I. M. Lifshits for useful advice and also V. G. Bar'yakhtar and S. V. Peletminskiy for profitable discussions. There are 9 references, 6 of which are Soviet.

ASSOCIATION: Fiziko-tehnicheskii institut Akademii nauk Ukrainsskoy SSR
(Physics and Technical Institute, AS UkrSSR)

SUBMITTED: March 22, 1958

Card 3/3

NOV 56-35-2-23/60

24161

AUTHORS:

Kaganov, M. I., Tsukernik, V. M.

TITLE:

The Influence of Thermoelectric Forces on the Skin Effect in Metal (Vliyaniye termoelektricheskikh sil na skan-effekt v metalle)

PERIODICAL:

Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1958, Vol 35, Nr 2, pp 474-478 (USSR)

ABSTRACT:

Formulae are deduced for the surface resistance in consideration of thermoelectric forces. The following system of equations serves as a basis:

$$\text{curl } \vec{H} = \frac{4}{c} \vec{j}; \text{curl } \vec{E} = -\frac{1}{c} \frac{d\vec{H}}{dt}; c \frac{\partial \Theta}{\partial t} + \text{div } \vec{q} = 0;$$

$$E_i = \rho_{ik} j_k + \alpha_{ik} \frac{\partial \Theta}{\partial x_k}; q_i = T \alpha_k j_k - \chi_{ik} \frac{\partial \Theta}{\partial x_k}$$

where Θ is the high-frequency addition to the average temperature T of the sample, c - the specific heat of the metal, \vec{q} the heat current, χ_{ik} - the resistance tensor, ρ_{ik} - the thermal conductivity tensor, and α_{ik} - the tensor of the thermoelectrical coefficients. ($\alpha_{ik} = \alpha_{ki}$).

Card 1/2

SCV/86-34-2-23/60

The Influence of Thermoelectric Forces on the Skin Effect in Metal

For the surface impedance it holds that $\xi_x = E_x(0)/H_y(0)$ and $\xi_y = -E_y(0)/H_x(0)$; $\xi_x = \xi_{0x} = \sqrt{\omega \sigma_{xx}/4\pi i} = \sqrt{\omega \sigma_{\perp}/4\pi i}$; ($\sigma_{\perp} = \sigma_{22} = \sigma_{33}$; $\sigma_{11} = \sigma_{11}$) and ξ_y is decomposed into ξ_y^{ad} and ξ_y^{isoth} , for which equations are deduced as well. In the case of isotropy $\xi = 1/\sqrt{\epsilon - \sin^2 \varphi}$ and $\xi^{isoth} = \sqrt{\epsilon - \sin^2 \varphi} / \epsilon$ is obtained, and for ξ^{ad} a considerably more complicated expression is found, the first term of which is equal to the expression for ξ^{isoth} . In conclusion, the authors express their gratitude to L. D. Landau for his valuable discussions. There are 2 references, 2 of which are Soviet.

ASSOCIATION: Fiziko-tehnicheskiy institut Akademii nauk ^USSR (Physico-Technical Institute, **AS UkrSSR**)

SUBMITTED: March 26, 1958
Card 2/2

TSUKERNIK, V.M.

24(0)
AUTHOR: Khatelnikov, I. M., Doctor of Physical and Mathematical Sciences
TITLE: Investigations of Low-temperature Physics (Issledovaniya po fizike nizkoi temperatur)

PERIODICAL: Vestnik Akademi nauk SSSR, 1959, Nr. 2, pp. 98-100 (USSR)
ABSTRACT: The 5th All-Union Conference on this problem took place in Tallin from October 27 to November 1, 1958. It was attended by physicists from Moscow, Kharkov, Leningrad, Tallin, Izdrevlensk, and Kiyev. 4 fields of low-temperature physics were discussed: superfluidity of liquid helium II, superconductivity, antiferromagnetism, magneto-resistive effect. The following reports and communications were heard: A. A. Abrikosov, G. I. Gor'khar reported on the investigation of the properties of superconductive alloys. A. A. Aronson, B. G. Bar'yakhtar, I. M. Khalilov, I. M. Khalilov reported on properties of superconductors in the high frequency magnetic field. D. M. Chiryaev and Chem. Chum'khan' reported on the investigation of the properties of superconductors. G. I. Gor'khar, I. M. Khalilov, and Chem. Chum'khan' described investigations for determination of the influence exercised by the Coulomb (Kulom) interaction of charges on superconductivity. V. V. Tolmachev explained the nature of the so-called collective excitations of the Bose type in superconductors. D. M. Zubary, Ya. A. Zaslavskiy spoke of the thermodynamics of superconductors and R. V. Geylman, V. Z. Kozlov of the thermal conduction of superconductors. In. V. Shadrin, V. F. Gornalshber reported on experimental work with superconductors. E. V. Kharzinskiy spoke of the measurement of the anisotropy of thermal conductivity in the superconductive state. In a series of reports problems of the superfluidity of helium were discussed, which was discovered in 1938 by P. L. Kapitza and the author, which was discovered in 1941 by P. L. Kapitza and the author, which was discovered by the author and the author. The properties of rotating helium. The interaction between superfluid and non superfluid helium. G. M. Yan'yan, collaborator of the Institut fizicheskikh problem (Institute of Physical Problems) investigated the properties of the so-called jump in temperature of Kapitza. I. M. Lifanil, V. M. Paschanskiy investigated galvanomagnetic phenomena in strong magnetic fields for metals with open Fermi surfaces. E. Ye. Alekseyevskiy, G. P. Gaydukov experimentally investigated the resistance anisotropy of gold microcrystals in the magnetic field. E. M. Kim, B. G. Bar'yakhtar, G. I. Gor'khar, A. temperature anisotropy with the accuracy of the theory of metallic conduction. I. M. Khalilov reported on the investigation of the properties of active field. A. A. Borovik-Borovoy reported on the weak ferromagnetism in antiferromagnetic samples of MnCO₃. E. M. Kopylov, I. M. Khalilov investigated the magnetic anisotropy of the antiferromagnetic monocrytals CuSO₄ and CoSO₄. E. A. Altkhanov reported on neutronographic investigations of antiferromagnetism. Ya. I. Kondoraki and collaborators reported on the susceptibility of nickel and nickel-copper alloys at low temperatures. M. I. Isenkov, I. M. Tsukernik reported on kinetic phenomena in ferromagnetics at low temperatures. A. I. Abrikosov, I. M. Khalilov, and S. P. Feinberg reported on the investigation of the magnetic anisotropy of ferromagnetic dielectrics at low temperatures. I. M. Khalilov reported on the results of paramagnetic resonance by terbins in the TMSO . 6H₂O nitrate. G. M. Khalilov gave a theoretical analysis of the effect in nonmetals. B. M. Smolyov, I. M. Kopylov and collaborators reported on obtaining orientated nuclei. E. V. Buktova, V. B. Kogan and J. C. Lasarevskiy reported on the investigation of the properties of the magnetic field in the presence of the magnetic field. I. M. Khalilov, V. B. Kogan, M. D. Starodubov and V. I. Koptevich detected polymorphisms in a number of metals at low temperatures. G. I. Aronkovich, V. P. Pashkov and M. P. Kalik reported on the stage of development of foreign scientific research work in the field of low-temperature physics. At the end of the Conference R. L. Kapliva spoke of his successful development of investigations in the field of low-temperature physics. The participants of the Conference visited the Institut fiziki Akademii nauk Gruzinskoy SSR (Physics Institute of the Academy of Sciences of the Gruzinskaya SSR) and the Physics Faculty of Tbilisi University as well as the building of the new research atomic reactor near Tbilisi.

Card 1/4

Card 2/4

90

24(3)

AUTHORS:

Kaganov, M. I., Tsukernik, V. M.

SOV/56-36-1-30/62

TITLE:

On the Phenomenological Theory of the Kinetic Processes
in Ferromagnetic Dielectrics (K fenomenologicheskoy teorii
kineticheskikh protsessov v ferromagnitnykh dielektrikakh)
II. The Interaction of Spin Waves With Phonons
(II. Vzaimodeystviye spinovykh voln s fononami)

PERIODICAL:

Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1959,
Vol 36, Nr 1, pp 224-232 (USSR)

ABSTRACT:

The spin waves have to be described by the vector of the
density of the magnetic moment $\vec{M} = \vec{M}(\vec{r}, t)$ by the deformation
tensor $u_{ik} = (1/2) (\partial u_i / \partial x_k + \partial u_k / \partial x_i)$ where $u = u(\vec{r}, t)$
denotes the vector of displacement. In this case, the interaction
Hamiltonian will be the sum of all components in the expansion
of the energy of the ferromagnetic with respect to the powers
of M_i , $\partial M_i / \partial x_k$, and u_{ik} : $\gamma_{iklm} \int M_i M_k u_{lm} dv$,

$\lambda_{iklmrs} = \int \frac{\partial M_i}{\partial x_k} \frac{\partial M_l}{\partial x_m} u_{rs} dv$. The first of these terms corresponds

Card 1/3

to the energy of magnetostriction. The interaction Hamiltonian

On the Phenomenological Theory of the Kinetic Processes in Ferromagnetic Dielectrics. II. The Interaction of Spin Waves With Phonons

SOV/56-36-1-30/52

is written down as $H_{\text{interaction}} = H_{\text{magnetostriction}} + H_{\text{exchange}}$,
 where $H_{\text{interaction}} = \gamma \int M_i M_k u_{ik} dv$,

$$H_{\text{exchange}} = \lambda_1 \int \frac{\partial M_1}{\partial x_i} \frac{\partial M_1}{\partial x_k} u_{ik} dv + \lambda_2 \int \frac{\partial M_1}{\partial x_i} \frac{\partial M_1}{\partial x_i} u_{kk} dv.$$

In the second part of the paper the production and the absorption of phonons by spin waves are calculated. The authors also investigate the energy transfer from spin waves to phonons. The spin waves and the phonons are described by equilibrium distribution functions of the Bose (Boze) type for different temperatures. The coefficient of heat transfer for the activation process of the emission and absorption of phonons decreases exponentially with temperature. The third part of the present paper deals with the annihilation of 2 spin waves with production of a phonon and with the decay of a phonon into 2 spin waves. These processes are due to the magnetostriction part of the Hamiltonian. Practically at any temperature above μMo , production and absorption of phonons

Card 2/3

On the Phenomenological Theory of the Kinetic Processes in Ferromagnetic Dielectrics. II. The Interaction of Spin Waves With Phonons

SOV/56-36-1-30/62

play the most important part among the one-phonon processes. The two-photon processes are described by the square terms of the expansion of the energy of the ferromagnetic with respect to the powers of the deformation tensor. There are 2 types of such processes: scattering of a phonon on a spin wave (or vice versa) and emission (or absorption) of 2 phonons by a spin wave. Finally, the time necessary for the establishment of equilibrium (relaxation time) between spin waves and phonons is calculated. There are 5 Soviet references.

ASSOCIATION: Fiziko-tehnicheskiy institut Akademii nauk Ukrainskoy SSR (Physico-Technical Institute of the Academy of Sciences, Ukrainskaya SSR)

SUBMITTED: July 8, 1958

Card 3/3

24 (3)
AUTHORS:

Kaganov, M. I., Tsukernik, V. M.

SOV/56-37-3-35/62

TITLE:

Nonresonance Absorption of the Energy of a Variable Magnetic Field by a Ferromagnetic Dielectrics

PERIODICAL:

Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1959,
Vol 37, Nr 3 (9), pp 823-832 (USSR)

ABSTRACT:

A ferromagnetic substance may absorb the energy of a variable magnetic field in various ways: either the magnetic field (the amplitude of which is to be small, $H' \sim e^{-i\omega t}$) is polarized perpendicular to the equilibrium magnetic moment, in which case the magnetic field turns the moment without changing its value, or the field is polarized in the direction of the moment, in which case a superposed magnetization occurs, i.e. the absolute value of the magnetic moment is varied. In both cases dissipative processes are caused, which are interrelated with the interaction of spin waves, both among themselves, and also with phonons (it is assumed that the dielectric substance is magnetized up to saturation). In the present paper the authors confine their attention to those in which the variable magnetic field (frequency ω) is polarized in the direction of the

Card 1/4

Nonresonance Absorption of the Energy of a Variable
Magnetic Field by Ferromagnetic Dielectrics

SOV/56-37-3-35/62

equilibrium moment. Calculation of absorption is then reduced to calculation of the imaginary part of longitudinal magnetic susceptibility $\mu_{||} = \mu'_{||} + i\mu''_{||}$. The complex character of the relaxation processes in ferromagnetic dielectrics leads to a complex frequency dependence $\mu''_{||}(\omega)$. The authors proceed from the simple model, in which the ferroelectric body is considered to have a magnetic moment, the vibrations of which propagate in form of spin waves. This is justified at sufficiently low temperatures, when the vibrations of one of the magnetic sublattices are not excited by the others. The theory of spin waves may be applied to real bodies, e.g. to ferrites the semiconductor properties of which at low temperatures are insignificant (the number of free electrons with decreasing temperature tends exponentially towards zero). The first part of this paper deals with the low frequency range, i.e. ω is much smaller than the reciprocal spin-spin relaxation time τ_{ss} : $\omega\tau_{ss} \ll 1$. The following is obtained for the imaginary part of the longitudinal magnetic susceptibility after all frequency- and temperature ranges have been dealt with in detail:

Card 2/4

Nonresonance Absorption of the Energy of a Variable
Magnetic Field by a Ferromagnetic Dielectrics

SOV/56-37-3-35/62

$$\mu'' \sim \begin{cases} \frac{M_0 \mu_i}{\theta_c} \left(\frac{T}{\theta_c}\right)^{1/2} \frac{\omega \tau_{sl}}{1 + \omega^2 \tau_{sl}^2}, & \omega \sqrt{\tau_{sl} \tau_{ss}} \ll 1, \quad T \ll T_0 \\ \frac{\mu M_0}{\theta_c} \left(\frac{T}{\theta_c}\right)^{1/2} \frac{\omega \tau_{ss}}{1 + \omega^2 \tau_{ss}^2}, & \omega \sqrt{\tau_{sl} \tau_{ss}} \gg 1, \quad T \ll T_0 \\ \frac{\hbar \omega}{\mu M_0} \frac{1 + \omega^2 \tau_2^2}{(1 + \omega^2 \tau_1^2)^2 + \omega^2 \tau_3^2}, & \omega \tau_{ss} \ll 1, \quad T \gg T_0 \end{cases}$$

Here $T_0 \sim \mu M_0 (\theta_c / \mu M_0)^{3/7}$, τ_{sl} denotes the spin-lattice relaxation time. The second part deals with high frequencies ($\omega \gg 1/\tau_{ss}$), the Hamiltonian of the interaction between field and spin being set up according to Kaganov (Ref 2):

$\mathcal{H}_{int} = \mu \int \hat{H}_z \hat{a}^*(\vec{r}) \hat{a}(\vec{r}) d\vec{r}$. If ω tends towards ∞ , then magnetic susceptibility tends towards unity, and its imaginary part towards zero. For the frequency dependence of μ'' at very high

Card 3/4

Nonresonance Absorption of the Energy of a Variable
Magnetic Field by a Ferromagnetic Dielectrics

SOV/56-37-3-35/62

frequencies it holds that $\mu'' \approx \frac{8\sqrt{2}}{15} \frac{\mu^2}{a^3} \frac{(2\pi\mu H_0)^2}{\theta_c^{3/2} (R\omega)^{3/2}}$,

$(R\omega \gg \frac{\lambda}{\tau_{SS}}, T, \mu H_{eff})$. Thus, μ'' decreases inversely

proportionally with $\omega^{3/2}$ and is independent of temperature.
The authors thank V. G. Ear'yakhtar for valuable discussions.
There are 10 Soviet references.

ASSOCIATION: Fiziko-tekhnicheskiy institut Akademii nauk Ukrainskoy SSR
(Physico-technical Institute of the Academy of Sciences,
Ukrainskaya SSR)

SUBMITTED: April 18, 1959

Card 4/4

В.М. Трубников

2(0)
AUTHOR:

Chantsov, R.

SOV/53-67-4-1/1

TITLE:

The Fifth All-Union Conference on the Physics of Low Temperature (5-ya Vsesoyuznoye soobshcheniye po fizike nizkikh temperatur)

SYNOPSIS:

Dapela fizicheskiy nauch. 1959, Vol 67, Nr 4, pp 743-750 (USSR)

ABSTRACT:

This conference took place from October 27 to November 1 at the Institute of Physics of the USSR Academy of Sciences. The main theme was the physics of low temperature. The conference was attended by about 100 specialists from the USSR and other countries. The conference was held in the USSR. About 50 lectures were delivered which were divided according to research fields.

III. Galvanomagnetic phenomena.

Card 5/11

(10 papers) I. R. Likhits and V. D. Paschenko (Khar'kov University) showed that the most important part in connection with the galvanomagnetic properties of metals is played by the concrete form of the Fermi surface of conductivity electrons. The author together with Yu. P. Gaidukov investigated the variation of the resistance in the framework of magnetic field with temperatures of Au, Cu, Pt, Fe, Ga, In and (Co)ZnO. V. I. Kovtina of St. Y. S. Borovik and (Co)ZnO. (KDFI) investigated the galvanomagnetic properties of the resistance of chromium group. A. Kan and N. G. Larinov (KDFI) investigated the resistance of the alloy in gold at low temperatures and found that the minimum of the resistance of the alloy in gold is connected with the temperature of the alloy in gold. The minimum of the resistance of the alloy in gold is connected with the temperature of the alloy in gold. The minimum of the resistance of the alloy in gold is connected with the temperature of the alloy in gold.

Card 6/11

Y. A. Alexandrov (KDFI) spoke about theoretical forces upon the skin effect in various conductors. Yu. V. Alexandrov (KDFI) spoke about the influence of the skin effect upon the skin effect in various conductors. Yu. V. Alexandrov (KDFI) spoke about the influence of the skin effect upon the skin effect in various conductors. Yu. V. Alexandrov (KDFI) spoke about the influence of the skin effect upon the skin effect in various conductors. Yu. V. Alexandrov (KDFI) spoke about the influence of the skin effect upon the skin effect in various conductors.

Card 7/11

Yu. A. Alexandrov (KDFI) spoke about his theory of the skin effect in various conductors. Yu. A. Alexandrov (KDFI) spoke about his theory of the skin effect in various conductors. Yu. A. Alexandrov (KDFI) spoke about his theory of the skin effect in various conductors. Yu. A. Alexandrov (KDFI) spoke about his theory of the skin effect in various conductors. Yu. A. Alexandrov (KDFI) spoke about his theory of the skin effect in various conductors.

V. M. TSOKERNIK

307/53-67-4-7/7

Chentsov, B.
The Fifth All-Union Conference on the Physics of Low
Temperatures (5-ye Vsesoyuznoye sobremennye po fizike nizkikh
temperatur)

24(O)
AUTHOR:

TITLE:

PERIODICAL:
(USSR)

ABSTRACT:
This Conference took place from October 27 to November 1 at
Tbilisi; it was organized by the Ordeleniye fiziko-matemati-
cheskikh nauk Akademii of SSSR (Department of Physico-
mathematical Sciences of the Academy of Sciences, USSR),
the Akademiya nauk Gruzinskoy SSR (Academy of Sciences,
Gruzinskaya SSR) and the Tbilisitskiy gosudarstvennyy uni-
versitet in S. Tbilisi (Tbilisi State University Lenin Collin).
The conference was attended by about 300 specialists from
Tbilisi, as well as by a number of young Chinese scientists
observing in the USSR. About 500 papers were deliv-
ered which were divided according to research fields.

Y. Nagatani.
A. S. Borovik-Romanov (IPP) delivered a report on investiga-
tions in carried out of the anisotropy of the weak ferro-
magnetism in monocrysal samples of the antiferromagnetic
MnCO (the effect of anisotropy was predicted by the thermo-
dynamic theory developed by Nagatani). In the course
of the discussion A. S. Borovik-Romanov (IPP) spoke about neutrono-
graphical investigations he carried out of the magnetic
structure of MnCO and FeCO at low temperatures. P. L.
Kaplan presented the importance of the method based upon
Pulchinskii's theory. I. K. Krasnaya (VNIIFRIL), whose
papers were read by A. S. Borovik-Romanov, reported on measur-
ments carried out by him (in the IPP) of the magnetic aniso-
tropy of the antiferromagnetic CuSO₄ and TlSO₄ monocrystals.

Card 7/11

Ye. A. Puzov (IPR AN SSSR, Sverdlovsk) spoke about his theo-
retical investigations of the magnetisability, the susceptibility,
the specific heat, and the resonance frequencies of anti-
ferromagnetics and ferrimagnetics. A. I. Rudovskiy and
Ye. A. Puzov (IPR AN SSSR) spoke about measurements of the
electric moments of iron in magnetic fields in a wide tem-
perature range with simultaneous plotting of the magnetization
curves. V. Tolikunskiy (IPR AN SSSR) spoke about measurements
of magnetization and the Hall effect of polycrystalline samples
nickel and Ni₂Mn at low temperatures. Ye. I. Kondratyuk

Y. Bode, B. Chan and Chang, Szu-chun (MCH) gave a report
on susceptibility measurements on nickel and its alloys with
copper at temperatures 2-10°K. S. S. Sandoz (TCU) spoke about
the anisotropy of the paramagnetic resonance of Tb³⁺ in terbium
halides at temperatures of liquid hydrogen. M. I. Kaganov
and Y. M. Tsokernik (MPTI) dealt with the kinetic phenomena
in ferromagnetics at low temperatures and with calculation
of relaxation times. A. I. Akhlyster, V. Sh. (VNIIFRIL) and
Ye. A. Puzov (IPR AN SSSR) carried out a theoretical investigation
of the relaxation of the magnetic moments in ferromagnetics;
Vlasov (IPR AN SSSR) showed that a highly polarized elastic
(ultrasonic) wave of a frequency of 10⁹ cycles when passing
through a ferromagnetic substance in the direction of the
magnetic field, is subjected to turn of the polarization
plane of the order of 10⁻³ - 10⁻⁴ rad/cm oriented. M. I.
Kaganov pointed out that in this case the resonance absorption
phenomenon may be observed, namely the resonance absorption
of ultrasonics if the wavelength is equal to the radius of
the Larmor orbit of the electron. Y. V. Vasilov, V. Vasilov,
and the most interesting facts.

Card 8/11

85974

S/126/60/010/005/029/030
E032/E414

24.2200 (1385, 1144, 1162)

AUTHORS: Kaganov, M.I., Tsukernik, V.M. and Chupis, I.Ye.
TITLE: Theory of Relaxation Processes in Antiferromagnetics ↗
PERIODICAL: Fizika metallov i metallovedeniye, 1960, Vol.10, No.5,
pp.797-798

TEXT: The method put forward by Akhiezer (Ref.1) and Kaganov and Tsukernik (Ref.2) is used to calculate the mean probabilities of processes associated with the interaction between spin waves in ferromagnetics. The theory holds in the temperature region

$$\omega_c \left(\frac{\mu_0 M_0}{\omega_c} \right)^{1/2} \ll T \ll \omega_c$$

✓

in which the energy of the spin wave is given by the well-known expression

$$\epsilon_\lambda = \omega_c (ak_\lambda)$$

Card 1/3

85974

S/126/60/010/005/029/030
E032/E414

Theory of Relaxation Processes in Antiferromagnetics

The notation is defined in the previous paper by the present authors (Ref.3). The processes considered are: (a) combination of two spin waves into one, and (b) collision of two spin waves. The probability of other processes, for example combination of three spin waves into one, have zero probability since energy and momentum conservation laws cannot be satisfied at the same time. It is found that the mean probabilities for the above two processes are respectively given by

$$\bar{W}_3 \approx \frac{\omega_c}{h} \left(\frac{\mu_0 M_0}{\omega_c} \right)^{5/2} \frac{T}{\omega_c}^3 \quad (3)$$

$$\bar{W}_4 \approx \frac{\omega_c}{h} \left(\frac{T}{\omega_c} \right)^5 \quad (4)$$

Comparison of these two probabilities shows that in the above
Card 2/3

85974

S/126/60/010/005/029/030
E032/E414

Theory of Relaxation Processes in Antiferromagnetics

temperature region the non-homogeneous exchange interaction is responsible for setting up the thermodynamic equilibrium in a spin wave system. Acknowledgments are expressed to V.G.Bar'yakhtara for valuable discussions. There are 3 Soviet references.

ASSOCIATIONS: Fiziko-tehnicheskiy institut AN USSR
(Physical-Technical Institute AS UkrSSR)
Khar'kovskiy gosudarstvennyy universitet
im. A.M.Gor'kogo (Khar'kov State University
im. A.M.Gor'kiy)

SUBMITTED: April 18, 1960

Card 3/3

83743

S/056/60/038/004/036/048
B006/B056

9.4300 (1035, 1138, 1143)
24.7900
AUTHORS:

Kaganov, M. I., Tsukernik, V. M.

TITLE:

Nonresonance Absorption of the Energy of an Alternating
Magnetic Field by a Ferromagnetic Dielectric. II

PERIODICAL:

Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1960,
Vol. 38, No. 4, pp. 1320 - 1325

TEXT: In part I of the present paper, the nonresonance absorption of the energy of an alternating magnetic field by a ferromagnetic dielectric has already been investigated on the assumption that this field is polarized in the direction of the axis of easiest magnetization. The case is now dealt with, in which the field is polarized perpendicular to this axis. These two cases are described as longitudinal and transverse fields. Contrary to what is the case in a longitudinal field, energy absorption may occur in a transverse field even if no dissipative processes occur, but when the field frequency agrees with the ferromagnetic resonance frequency. Neither resonance nor the effects connected with it are considered. The results obtained here relate to frequencies

X

Card 1/3

83743

Nonresonance Absorption of the Energy of an Alternating Magnetic Field by a Ferromagnetic Dielectric. II

S/056/60/038/004/036/048
B006/B056

that are distant from resonance. The dissipative processes connected with the energy absorption are caused by the interaction between the magnetic field and the spin waves. The authors assume (like in Ref. 1) that the ferromagnetic dielectric is magnetized up to saturation at a given temperature, i.e., that it may be considered to be a single-domain sample. The sample is also considered to be sufficiently pure, so that impurity effects may be neglected. It is shown that the nonresonance absorption of magnetic field energy is connected with the forming of two spin waves as a result of a "collision" between a photon and a spin wave. Besides, an absorption of the photon in higher perturbation-theoretical approximations is possible due to processes in which a large number of spin waves participate. The most important process in this case is one of the fifth order, which occurs in second perturbation-theoretical approximation. The coefficient of the absorption of magnetic field energy $\Gamma = Q \left(\frac{h\omega}{8\pi} V \right)^{-1}$ is now investigated, where Q denotes the amount of energy absorbed per second within the entire sample volume.

Card 2/3

83743

Nonresonance Absorption of the Energy of an Alternating Magnetic Field by a Ferromagnetic Dielectric. II

S/056/60/038/004/036/048
B006/B056

One obtains $\Gamma = \frac{1}{32\pi^2} \frac{w^2 T^2}{\mu M_0 \theta_c^3} \omega (1 - e^{-\hbar\omega/T}) I(\eta, \nu)$, where $\eta = \mu H_e/T$ and

$\nu = \hbar\omega/T$; the function $I(\eta, \nu)$ is represented as an integral equation by (13) and (13'). These equations are integrated for the special cases of low frequencies ($\nu \ll 1, \eta$), resonance frequency ($\nu = \eta$), and high frequencies ($1 \gg \nu \gg \eta$; $\nu \gg 1 \gg \eta$; $\eta \gg \nu \gg 1$; $\nu \gg \eta \gg 1$). The resulting asymptotic values of the integrals are used to determine approximate formulas for Γ in the various special cases. The authors thank A. I. Akhiezer and V. G. Bar'yakhtar for discussions. There are 3 references: 2 Soviet and 1 US.

ASSOCIATION: Fiziko-tekhnicheskiy institut Akademii nauk Ukrainskoy SSR
(Institute of Physics and Technology of the Academy of
Sciences Ukrainskaya SSR)

SUBMITTED: November 23, 1959

Card 3/3

S/056/60/039/002/044/044
B006/B070

AUTHORS: Kaganov, M. I., Tsukernik, V. M.

TITLE: Theory of the Non-resonant Absorption of a Variable
Magnetic Field by a Ferromagnetic Dielectric

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1960,
Vol. 39, No. 2(8), p. 518

TEXT: In an earlier paper, the authors calculated the absorption coefficient of a variable magnetic field polarized perpendicular to the axis of easiest magnetization. The Hamiltonian used in that case took only the exchange interaction into account. Later, Tsukernik showed that in this case only resonance absorption of a homogeneous magnetic alternating field may take place, since the total magnetic moment commutes with the Hamiltonian of the system; the results of the earlier paper (Ref. 1) are therefore wrong. The error is related to the fact that in the calculation of the matrix elements of the transition, the authors restricted themselves to the first approximation of the perturbation theory. A consideration of the second approximation shows that the matrix

Card 1/2

Theory of the Non-resonant Absorption of a
Variable Magnetic Field by a Ferromagnetic
Dielectric

S/056/60/039/002/044/044
B006/B070

element vanishes to the corresponding accuracy. The non-resonant absorption of the homogeneous field is related to the relativistic interactions in the system (dipole-dipole interaction, anisotropy energy, etc.). These problems will be considered in a later paper. Here, the authors give only a formula for the coefficient of absorption of a transverse magnetic field whose frequency is considerably smaller than that of the spin wave in the case of a zero quasi-momentum. The absorption described by this formula is due to a decay of a photon into two spin waves with oppositely directed quasi-momenta. There is 1 Soviet reference. ✓

ASSOCIATION: Fiziko-tehnicheskii institut Akademii nauk Ukrainskoy SSR
(Institute of Physics and Technology of the Academy of
Sciences Ukrainskgya SSR)

SUBMITTED: July 2, 1960

Card 2/2

ACCESSION NR: APL019844

S/0181/64/006/003/0818/0826

AUTHORS: Genkin, G. M.; Golubeva, N. G.; Tsukornik, V. M.

TITLE: The spin phonon width of lines of antiferromagnetic resonance

SOURCE: Fizika tverdogo tela, v. 6, no. 3, 1964, 818-826

TOPIC TAGS: spin phonon interaction, magnetic resonance, magnetization precession, exchange interaction, antiferromagnetism

ABSTRACT: The authors have examined the relaxation of the uniform precession of magnetization in a uniaxial antiferromagnetic as a consequence of interaction between spin waves and phonons (both acoustical and optical). They show that the relaxation processes of creating (or absorbing) an acoustical phonon by a spin wave, by virtue of the law of conservation of energy and momentum, may take place only at frequencies of

$$\omega_0 > \omega_0 \pm \frac{\sqrt{3}h_A v}{a}$$

where ω_0 is the frequency of antiferromagnetic resonance, h_A is the field of

Card 1/2

ACCESSION NR: AP4019844

anisotropy, v the velocity of sound, and a the lattice constant. The interaction is an exchange process. The width of the line has been computed for any temperature. At rather high frequencies of antiferromagnetic resonance, $\omega_0 > \omega_0^{ex}$, where normally $\omega_0^{ex} \sim 2 \cdot 10^{12}$ sec⁻¹deg, the spin-phonon line width at low temperatures proves to be much greater than spin-spin line width. At frequencies of $\omega_0 < \omega_0^{ex}$, the spin-phonon width is small compared to spin-spin width, except for regions of very low temperatures ($T < 10^{-3}$ K). Orig. art. has: 39 formulas.

ASSOCIATION: Radiofizicheskiy institut, Gorkiy (Radiophysical Institute)

SUBMITTED: 21Sep63

DATE ACQ: 31Mar64

ENCL: 00

SUB CODE: SS, EM

NO REF SOV: 003

OTHER: 008

Card 2/2

24.7900

1147 1158 1163, 1144

30083
S/048/61/025/C11/007/031
B108/B138

AUTHORS: Kaganov, M. I., and Tsukernik, V. M.

TITLE: Off-resonance absorption of a variable magnetic field by ferromagnetic dielectrics at low temperatures

PERIODICAL: Akademiya nauk SSSR. Izvestiya. Seriya fizicheskaya, v. 25, no. 11, 1961, 1346-1351

TEXT: The authors studied the absorption of an h.f. magnetic field whose plane of circular polarization is perpendicular to the equilibrium magnetic moment of the ferromagnetic. They consider absorption without resonance, which can only occur by the interaction of spin waves with one another and with phonons. The calculations are made for temperatures considerably below the Curie point θ_c . The absorption coefficient Γ of the magnetic field is given as $\Gamma = \frac{8\pi k^2 \omega}{h^2 V} \sum \{w_{if} - w_{fi}\}$, where w_{if} is the probability of direct, and w_{fi} of inverse, transition. The absorption

X

Card 1/4

30063 S/048/61/025/011/007/031
B108/B138

Off-resonance absorption of a...

coefficient due to spin-spin interaction is evaluated for two limiting cases: (1) High temperatures $T \gg \epsilon_0$ (but still $T \ll \theta_c$):

$$\Gamma_{ss} \approx \begin{cases} A_1 \omega \left(\frac{w}{\theta_c}\right)^2 \frac{\mu M_1}{\theta_c} \cdot \frac{\hbar \omega T^2}{\epsilon_0^3}, & \hbar \omega \ll \epsilon_0, \\ A_2 g M_0 \left(\frac{w}{\theta_c}\right)^2 \frac{T^2}{\hbar \omega \theta_c}, & \epsilon_0 \ll \hbar \omega \ll T, \\ \frac{4}{15} \sqrt{\frac{\pi}{2}} \zeta(3/2) g M_0 \left(\frac{w}{\theta_c}\right)^2 \frac{T^{3/2}}{(\hbar \omega)^{1/2} \theta_c}, & \hbar \omega \gg T, \end{cases} \quad (17)$$

where $w = \frac{\mu^2}{a^3}$ is the dipole-dipole interaction energy; $A_1 \sim A_2 \sim 10^3$.

(2) Low temperatures ($T \ll \epsilon_0$):

Card 2/4

30053 3/048/61/025/011/007/031
B100/B136

Off-resonance absorption of a...

$$\Gamma_{33} \approx \begin{cases} B_1 \omega \left(\frac{w}{\theta_c}\right)^2 \frac{\mu M_0}{\theta_c} \frac{\hbar \omega T^{1/2}}{\epsilon_0^{1/2}} e^{-\frac{3t_a}{T}}, & \hbar \omega \ll \epsilon_0, \\ \frac{4}{15} \sqrt{\frac{\pi}{2}} \omega \left(\frac{w}{\theta_c}\right)^2 \frac{\mu M_0}{\theta_c} \left(\frac{T}{\epsilon_0}\right)^{1/2} e^{-\frac{3t_a}{T}}, & \hbar \omega \gg \epsilon_0, \end{cases} \quad (18)$$

where

$$B_1 = \frac{15\pi^{1/2}}{8\sqrt{2}} \left\{ 121 + \frac{1}{15} \left[\left(\frac{2\beta}{\pi} - \frac{2}{3}\right)^2 + \frac{121}{45} \right] \right\} + \frac{4}{15} \sqrt{\frac{\pi}{2}} \approx 10^4$$

$\epsilon_0 = \mu H_0 + \beta \mu M_0$ where H_0 denotes the permanent magnetic field along the axis of easiest magnetization, M_0 the equilibrium magnetic moment, β the anisotropy constant, a the lattice constant, $\mu = g\hbar$, g the gyromagnetic ratio. The results for spin-lattice interaction show that these contributions to the overall absorption coefficient are considerably smaller than the contribution due to spin-spin interaction. The considerations have been made for perfect ferromagnetics without

X

Card 3/4

Off-resonance absorption of a...

3/046/61/025/011/007/031
B108/B138

impurities. I. M. Lifshits is thanked for discussions. This paper was read at the Conference on ferromagnetism and antiferromagnetism in Leningrad, May 5-11, 1961. There are 1 figure and 4 Soviet references.

ASSOCIATION: Fiziko-tekhnicheskiy institut Akademii nauk USSR
(Physicotechnical Institute of the Academy of Sciences
UkrSSR)

Card 4/4

24.7900 (1055, 1144, 1163)
24.2200 1164 1068 1121

30064
S/048/61/025/011/008/031
B108/B138

AUTHORS: Kaganov, M. I., and Tsukernik, V. M.

TITLE: An absorption mechanism for a longitudinal magnetic field by a ferromagnetic dielectric

PERIODICAL: Akademiya nauk SSSR. Izvestiya. Seriya fizicheskaya, v. 25, no. 11, 1961, 1352-1353

TEXT: The ferro-dielectric absorption of the energy of a variable magnetic field polarized along the axis of easiest magnetization was studied in detail in a previous paper (Ref. 1: Zh. eksperim. i teor. fiz., 37, 823 (1959)). The absorption of the field at $\omega\tau_{ss} \ll 1$, where τ_{ss} is relaxation time, was calculated with the aid of the kinetic equations for the distribution function of the spin waves. The calculations were made for temperatures T considerably below Curie point θ_c . At high frequencies, absorption is chiefly determined by the disintegration of a photon into two spin waves with opposite momenta. This process was also considered in the above paper. It was pointed out, however, that a photon will only

Card 1/4

30064
S/048/61/025/011/008/031
B108/B138

An absorption mechanism for a...

disintegrate into two spin waves at frequencies ω greater than $\frac{2\epsilon_0}{\hbar}$, where ϵ_0 is the energy of a zero-momentum spin wave. The wide range $\frac{1}{\tau} \ll \omega < \frac{2\epsilon_0}{\hbar}$ has not been considered as yet. Absorption in this range is the subject of the present paper. In the lowest spin-wave approximation, the Hamiltonian for spin wave interaction with one another and with a longitudinal magnetic field is $H' = H_{int} + H_h$ (1) where $H_{int} = \sum_{123} \Phi_{12;3} a_1^+ a_2^+ a_3 + \text{complex conjugate}$ (2) and $H_h = \frac{1}{2} \mu h_0 (e^{i\omega t} + e^{-i\omega t}) \sum_k a_k^+ a_k$ (2'). h_0 denotes the amplitude of

the variable field. The other notations may be found in Ref. 2 (Akhiezer A.I. et al., Uspekhi fiz. nauk, 71, 533 (1960)). In the approximation in question (second approximation in perturbation theory), strong exchange interaction does not contribute to absorption. It can be seen from the (1) and (2) that the transition matrix elements that are linear with respect to the variable field are nonvanishing only in second and higher perturbation-theoretical approximation. The corresponding lowest order processes are: (a) merging of a photon and a spin wave, which form two spin waves and (b) merging of two spin waves and one photon to form one spin wave. The

Card 2/4

30064 S/048/61/025/011/008/031
B108/B138

An absorption mechanism for a...

calculation of the appropriate absorption coefficients Γ_a and Γ_b is similar to that given in detail in Ref. 3 (Kaganov M. I., Tsukernik V. M., present number of this periodical, p. 1346). In this article, only the final results for the total absorption coefficient $\Gamma = \Gamma_a + \Gamma_b$ are given, with precision up to a numerical factor of the order of unity. At high temperatures ($\epsilon_0 \ll T \ll \theta_c$)

$$\Gamma \sim \begin{cases} gM_0 \frac{w^3}{\epsilon_0 \theta_c} \left(\frac{T}{\theta_c}\right)^2, & \frac{\hbar}{\tau_{sp}} \ll \hbar\omega \ll \epsilon_0, \\ gM_0 \frac{w^3}{\hbar\omega \theta_c} \left(\frac{T}{\theta_c}\right)^2, & \epsilon_0 \ll \hbar\omega \ll T, \\ gM_0 \frac{w^3}{\theta_c^{3/2} (\hbar\omega)^{1/2}} \left(\frac{T}{\theta_c}\right)^{3/2}, & T \ll \hbar\omega \ll \theta_c. \end{cases}$$

At low temperatures ($T \ll \epsilon_0$)

Card 3/4

An absorption mechanism for a...

30064 S/048/61/025/011/008/031
B108/B138

$$\Gamma \sim \begin{cases} gM_0 \frac{w^2}{\theta_c^2} \left(\frac{\epsilon_0}{\theta_c}\right)^{1/2} \left(\frac{T}{\theta_c}\right)^{1/2} e^{-\frac{3\epsilon_0}{T}}, & \frac{\hbar}{\tau_{11}} \ll \hbar\omega \ll \epsilon_0, \\ gM_0 \frac{w^2}{\theta_c^{1/2} (\hbar\omega)^{1/2}} \left(\frac{T}{\theta_c}\right)^{1/2} e^{-\frac{\epsilon_0}{T}}, & \hbar\omega \gg \epsilon_0. \end{cases}$$

X

Comparison of the results obtained with the absorption coefficient due to the disintegration of one photon into two spin waves (Ref. 1) shows that the processes considered in this paper are only effective at $\hbar\omega < 2\epsilon_0$.
 [Abstracter's note: Complete translation.] This paper was read at the Conference on ferromagnetism and antiferromagnetism in Leningrad, May 5-11, 1961. There are 3 Soviet references.

ASSOCIATION: Fiziko-tekhnicheskii institut Akademii nauk USSR
 (Physicotechnical Institute of the Academy of Sciences
 UkrSSR)

Card 4/4

26) 21
S/056/61/041/001/019/021
B102/B231

24.2200

AUTHORS: Kaganov, M. I., Tsukernik, V. M.

TITLE: High-frequency magnetic susceptibility of uniaxial ferromagnetic crystal in a longitudinal magnetic field

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, v. 41, no. 1(7), 1961, 267 - 271

TEXT: Susceptibility dispersion of an antiferromagnetic material is mainly due to rotation of the magnetic moments of the sublattices in a magnetic field. The imaginary part of χ exhibits a resonance character in this case: resonance frequency coincides with the natural rotation frequencies of the system of moments. These frequencies have been computed for various equilibrium configurations by C. Kittel (Phys. Rev., 85, 329, 1952) and Ye. A. Turov (ZhETF, 34, 1009, 1958). In an earlier paper (ZhETF, 24, 524, 1958), the authors of the present work have computed the h-f susceptibility of a uniaxial antiferromagnetic body without magnetic field. If the magnetic field is strong enough, the equilibrium configuration of the sublattice moments may change, which

Card 1/7

25
X

26424

S/056/61/041/001/019/021

B102/B231

High-frequency magnetic ...

would result in a change of the dispersion character. Starting from the Landau-Lifshits equations for the motion of the sublattice moments, the h-f susceptibility tensor of an antiferromagnetic body is computed for various values of the constant magnetic field strength, that is, for various equilibrium configurations of the moments. The results permit to draw conclusions from h-f measurements as to the equilibrium structures and the transitions between them. The uniaxial antiferromagnetic body is assumed to have two sublattices and to be positioned in a homogeneous, constant and in a weak and variable (frequency ω) magnetic field. The motion of the moments due to field action is described by

$$\partial M_s / \partial t = g [M_s H_e^{(s)}] - (\gamma / M_s^2) [\vec{M}_s [M_s H_e^{(s)}]] \quad (1)$$

where \vec{M}_s is the magnetization vector of the s-th sublattice, M the sublattice magnetization which is assumed to be constant, g the gyromagnetic ratio, γ the relaxation constant, $H_e^{(s)}$ the effective field acting upon the s-th sublattice: $H_e^{(s)} = -\partial \chi / \partial \vec{M}_s$, where χ stands for the energy density of the antiferromagnetic body:

Card 2/7

High-frequency magnetic ...

26424
S/056/61/041/001/019/021
B102/B231

$$\mathcal{H} = \alpha M_1 M_2 - \frac{1}{2} \lambda [(M_1 n)^2 + (M_2 n)^2] + \eta (M_1 n) (M_2 n) - H (M_1 + M_2). \quad (3)$$

Here, α is the constant of exchange interaction ($\alpha > 0$), λ and η represent the anisotropy constants which are assumed to be positive; n is the unit vector in the direction of the axis of the antiferromagnetic body. Using the notation stated hereinafter: $H_1 = \sqrt{(\lambda + \eta)(2\alpha - \lambda + \eta)} M$,

$$H_2 = \sqrt{(\lambda + \eta)(2\alpha + \lambda + \eta)} M, \quad H = H_3 = (2\alpha - \lambda + \eta) M, \quad \text{and}$$

$$H_4 = [\lambda(2\alpha - \lambda + \eta)^2 / (2\alpha + \lambda + \eta)]^{1/2} M \quad (H_4 < H_1; H_2 - H_4 \text{ is the width of}$$

the hysteresis loop of the antiferromagnetic body), the following cases are subjected to investigation: 1) $H < H_1$. m_{\pm} is equal to $\chi_{\pm} h_{\pm}$, where

$h_{\pm} = h_x \pm i h_y$, $m_{\pm} = m_x \pm i m_y$, \vec{h} stands for the h-f magnetic field, and \vec{m} for the variable part of the entire magnetic moment. With

$$\Omega^2 = (g^2 M^2 + \gamma^2) (\lambda + \eta) (2\alpha + \lambda + \eta) - \gamma^2 H^2 / M^2, \quad (5)$$

$$\Omega_1^2 = 2 (g^2 M^2 + \gamma^2) (\lambda + \eta),$$

Card 3/7

26424
S/056/61/041/001/019/021
B102/B231

High-frequency magnetic ...

the following is obtained:

$$\chi_{\pm} = \frac{\Omega_{\pm}^2 - 2i\omega\gamma}{\Omega^2 - (\omega \mp gH)^2 - 2i\omega\gamma(\alpha + \lambda + \eta)} \quad \chi_{zz} = 0. \quad (6)$$

The width of the antiferromagnetic resonance lines is thus given by $2\gamma(\alpha + \lambda + \eta)$. With $H = 0$ one has $\chi_{+} = \chi_{-} = \chi_{xx} = \chi_{yy}$. 2) $H_1 < H < H_3$; Here, $\chi_{zz} \neq 0$, and the following is obtained:

$$\chi_{zz}(\omega) = \chi_{zz}(0) \frac{v^2 + i\nu\omega}{v^2 + \omega^2}; \quad (7)$$

$$\chi_{zz}(0) = 2/(2\alpha - \lambda + \eta). \quad (8)$$

$$v = (2\alpha - \lambda + \eta) \gamma \sin^2 \theta = \gamma (1 - H^2/H_3^2) H_3/M.$$

From this, the following is obtained for the behavior of the relaxation time $\tau = 1/\nu$ in the vicinity of the point of second-kind phase transition (with $H \approx H_3$):

$$\tau = \frac{1}{\delta} \frac{H_3}{H_3^2 - H^2}. \quad \text{If } T \text{ is fixed, and } H = H_3 \text{ the following is valid:}$$

Card 4/7

26424
S/056/61/041/001/019/021
B102/B231

High-frequency magnetic ...

$$\tau = \frac{G}{|T_k - T|}, \quad G = M/2\gamma \left| \frac{dH_s}{dT} \right|_{T=T_k} \quad (8'')$$

where T_k is the temperature of phase transition. This means that with the frequency being fixed there is a field or a temperature at which the h-f field (polarized along z) exhibits an absorption maximum whose position and height is dependent on ω . The transverse components are in this case given by

$$\begin{aligned} \chi_{xx} &= \frac{\omega_1^2 - 2i\omega\gamma}{\omega_0^2 - \omega^2 - 2i\alpha\gamma'\omega} \cos^2 \theta, \\ \chi_{yy} &= \frac{1}{\alpha} \frac{\omega_0^2 - 2i\omega\gamma}{\omega_0^2 - \omega^2 - 2i\alpha\gamma'\omega}, \\ \chi_{xy} &= -\chi_{yx} = \frac{2igM\omega \cos \theta}{\omega_0^2 - \omega^2 - 2i\alpha\gamma'\omega} \end{aligned} \quad (9)$$



Card 5/7

26: 2h

S/056/61/041/001/019/021
B102/B231



High-frequency magnetic ...

where

$$\begin{aligned} \omega_0^2 &= (g^2 M^2 + \gamma^2) (4\alpha^2 \cos^2 \theta - 2(\lambda + \eta) \alpha \sin^2 \theta), \\ \omega_1^2 &= 4\alpha (g^2 M^2 + \gamma^2), \quad \gamma' = \gamma \left[1 + \cos^2 \theta + \frac{1}{2\alpha} (\lambda + \eta) \sin^2 \theta \right]. \end{aligned} \quad (10)$$

3) $H > H_3$, ($\vec{H} \parallel \vec{n}$). Here, the tensor of h-f susceptibility coincides with that of the susceptibility of the uniaxial antiferromagnetic body:

$$\begin{aligned} \chi_{xx}(\omega) = \chi_{yy}(\omega) = \chi_{zz}(\omega) &= \chi_{\perp}(0) \frac{\omega_{\phi}^2 - i\omega\gamma_{\phi}}{\omega_{\phi}^2 - \omega^2 - 2i\omega\gamma_{\phi}}, \\ \chi_{xy}(\omega) = -\chi_{yx}(\omega) &= \frac{2igM\omega}{\omega_{\phi}^2 - \omega^2 - 2i\omega\gamma_{\phi}}, \end{aligned} \quad (11),$$

$$\chi_{xz} = \chi_{yz} = \chi_{zx} = 0,$$

$$\begin{aligned} \chi_{\perp}(0) &= 2M/(H + (\lambda - \eta)M), \\ \omega_{\phi}^2 &= g^2 [H + (\lambda - \eta)M]^2 (1 + \gamma^2/g^2 M^2), \\ \gamma_{\phi} &= \gamma (H/M + \lambda - \eta). \end{aligned} \quad (12).$$

Card 6/7

High-frequency magnetic ...

26424
S/056/61/041/001/019/021
B102/B231

With $H = H_3 = (2\alpha - \lambda + \eta)M$ all components of $\chi_{ik}(\omega)$ are continuous.
The authors thank A. S. Borovik-Romanov for remarks. There are
7 references: 3 Soviet-bloc and 4 non-Soviet-bloc.

ASSOCIATION: Fiziko-tehnicheskii institut Akademii nauk Ukrainskoy SSR
(Institute of Physics and Technology of the Academy of
Sciences Ukrainskaya SSR)

SUBMITTED: February 17, 1961

X

Card 7/7

R0227

S/056/62/043/003/037/063
B108/B102

24.11.00

AUTHORS: Gurzhi, R. N., Tsukernik, V. M.

TITLE: High-frequency magnetic susceptibility of ferromagnetic dielectrics in a longitudinal magnetic field

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, v. 43, no. 3(9), 1962, 977 - 983

TEXT: Using a method-similar to Bogolyubov's (ZhETF, 17, 614, 1947) the authors got the following quantum kinetic equation (linear approximation with regard to the variable field $h_t = h_0 e^{-i\omega t}$) for the distribution function $F_k = \langle a_k^+ a_k \rangle$ of spin waves:

$$-i\omega f_k = L_k h_0 + \hat{W} f_k; \tag{11}$$

$$L_k = (\pi\mu\omega/2) |B_k|^2 (2\tilde{n}_k + 1) \tilde{e}_k^{-2} [\delta_-(-2\tilde{e}_k - \hbar\omega) - \delta_-(2\tilde{e}_k - \hbar\omega)] +$$

$$+ 4\pi\mu\omega \sum_{k_1} \frac{|\Phi_{k_1, 2}|^2}{e_{k_1, 2}^2} [(1 + n_k + n_1) n_2 - n_1 n_k] [\delta_-(e_{2, 1k} - \hbar\omega) - \delta_-(e_{k_1, 2} - \hbar\omega)] +$$

Card 1/4

High-frequency magnetic...

S/056/62/043/003/037/063
B108/B102

$$\begin{aligned}
 & + 2\pi i \omega \sum_{12} \frac{|\Phi_{12;k}|^2}{e_{12;k}^2} [(1 + n_1 + n_2)n_k - n_1 n_2] [\delta_-(e_{12;k} - \hbar\omega) - \delta_-(e_{k;12} - \hbar\omega)], \\
 \mathcal{W}f_k = & \frac{4\pi}{\hbar} \sum_{12} |\Phi_{k1;2}|^2 [(1 + n_k + n_1) f_2 + (n_2 - n_k) f_1 + (n_2 - n_1) f_k] \times \\
 & \times [\delta_-(e_{k1;2} - \hbar\omega) + \delta_-(e_{2;k1} - \hbar\omega)] + \frac{2\pi}{\hbar} \sum_{12} |\Phi_{12;k}|^2 [(n_2 - n_k) f_1 + \\
 & + (n_1 - n_k) f_2 - (1 + n_1 + n_2) f_k] [\delta_-(e_{k;12} - \hbar\omega) + \delta_-(e_{12;k} - \hbar\omega)] + \\
 & + \frac{16\pi}{\hbar} \sum_{123} |\Phi_{k1;23}|^2 \{ [n_1(n_2 + 1)(n_3 + 1) - (n_1 + 1)n_2 n_3] f_k + [n_k(n_2 + 1) \times \\
 & \times (n_3 + 1) - (n_k + 1)n_2 n_3] f_1 + [n_k n_1(n_3 + 1) - (n_k + 1)(n_1 + 1)n_3] f_2 + \\
 & + [n_k n_1(n_2 + 1) - (n_k + 1)(n_1 + 1)n_2] f_3 \} [\delta_-(e_{k1;23} - \hbar\omega) + \\
 & + \delta_-(e_{23;k1} - \hbar\omega)].
 \end{aligned}$$

Card 2/4

S/056/62/043/003/037/063
B108/B102

High-frequency magnetic...

with $F_\alpha|_{t=0} = n_\alpha = (e^{\epsilon_\alpha/T} - 1)^{-1}$ and $\langle a_\alpha^+ a_\beta^+ a_\gamma \rangle|_{t=0} = 0$.

$\epsilon_k = \theta_{\text{Curie}}(ak)^2 + \mu H + |B_k|$. At high frequencies it is sufficient to consider the term $-i\omega f_k = L_k h_0$. From this equation the high-frequency magnetic susceptibility of a ferroelectric with one magnetic sublattice can be calculated keeping in mind that magnetization too can be expressed in terms of the spin wave distribution function:

$$V^{-1} \langle M_z \rangle = M_0 - \frac{\mu}{V} \sum_k \langle a_k^+ a_k \rangle, \quad \langle a_k^+ a_k \rangle = \text{Sp } \rho a_k^+ a_k, \quad (6)$$

Thus, $\chi = -\frac{i\mu V}{\omega V} \sum_k L_k$. V is the volume of the body. The real part of magnetic susceptibility which is related to dispersion is

$$\text{Re } \chi = \frac{\mu}{\omega V} \sum_k \text{Im } L_k$$

The predominant term in $\text{Im } L_k$ is the one proportional to $|B_k|^2$. Con-

Card 3/4

High-frequency magnetic...

S/056/62/043/003/037/063
B108/B102

sidering this and using the symbols $x = \sqrt{\theta_{\text{Curie}}/T} ak$, $\alpha = \hbar\omega/T$,
 $\eta = \mu H/T$, $\gamma = 4\pi\mu M_0/T$ and $w = \mu^2/a^3$ the result is

$$\text{Re}\chi = \frac{w \cdot (\mu M_0)^2}{4T} \left(\frac{T}{\theta_c}\right)^{3/2} \int_0^\pi \sin^3 \theta d\theta \int_{-\infty}^{\infty} \frac{\text{ch}(y(x)/2)}{y(x)} \frac{x^2 dx}{y^2(x) - a^2/4}$$

$$y(x) = \sqrt{(x^2 + \eta)^2 + \gamma(x^2 + \eta) \sin^2 \theta}$$

ASSOCIATION: Fiziko-tehnicheskii institut Akademii nauk Ukrainsskoy SSR
(Physicotechnical Institute of the Academy of Sciences of the
Ukrainsskaya SSR)

SUBMITTED: March 31, 1962

Card 4/4

L 40173-66 EWT(1) JW

ACC NR: AP6018817 SOURCE CODE: UR/0056/66/050/005/1377/1380

AUTHOR: Pikin, S. A.; Tsukernik, V. M.

ORG: none

TITLE: Thermodynamics of linear spin chains in a transverse magnetic field

SOURCE: Zh eksper i teor fiz, v. 50, no. 5, 1966, 1377-1380

TOPIC TAGS: spin system, transverse magnetic field, magnetic susceptibility, THERMODYNAMIC PROPERTY, HOMOGENEOUS MAGNETIC FIELD

ABSTRACT: A one-dimensional system of spins ($s = 1/2$) with a strongly anisotropic interaction of the immediate neighbors has been examined. The system is assumed to be in a homogeneous magnetic field directed along a given axis. Exact equations have been obtained for the thermodynamic characteristics of the system. It has been shown that at $T = 0$ and an absolute value of the magnetic-field strength H_0 , the magnetic susceptibility has a logarithmic or root nature, dependent, like H_0 , on the relation between the interaction constants. The authors thank V. G. Vaks and A. I. Larkin for valuable discussions. Orig. art. has: 11 formulas. [Based on authors' abstract] [NT]

SUB CODE: 20/ SUBM DATE: 03Dec65/ ORIG REF: 003/ OTH REF: 002

L 41744-66 EWT(1) IJP(c) GG

ACC NR: AF6020219

SOURCE CODE: UR/0056/66/050/006/1631/1635

AUTHOR: Tsukernik, V. M.

61

ORG: none

B

TITLE: Features of gyromagnetic effect in ferroelectrics at low temperatures

SOURCE: Zh eksper i teor fiz, v. 50, no. 6, 1966, 1631-1635

TOPIC TAGS: ferromagnetic dielectric, spin wave theory, dipole interaction, magnetic moment, ground state, temperature dependence

ABSTRACT: The author derives in the spin-wave approximation a formula for the temperature part of the mechanical moment of a spin system and shows that in the region of temperatures where the dipole interaction plays an important role, its ratio to the corresponding addition to the magnetic moment differs from the single-electron g factor and depends on the temperature. The approximation employed consists of expanding, with accuracy to quadratic terms, the energy density in powers of a quantity which is small near the ground state in comparison with the magnetization, followed by quantization and averaging with an equilibrium Gibbs distribution. The deduced difference in the temperature dependence of the magnetic and mechanical moments of the spin system is observable in principle with the aid of a gyromagnetic experiment in which the moments are changed as a result of changes in the temperature in the absence of an external field. The author thanks M. I. Kaganov for useful discussions. Orig. art. has: 20 formulas.

SUB CODE: 20/ SUBM DATE: 18Jan65/ ORIG-REF: 004

Card 1/1

SOURCE: Zhurnal eksperimental'noy i teoreticheskoy fiziki,
v. 48, no. 5, 1965, 1386-1392

TOPIC TAGS: electromagnetic field, field state, field intensity
value, wave function, occupation number, quantum electrodynamics

ABSTRACT: Since a free electromagnetic field can be described in
quantum theory either by giving the photon occupation numbers or
by giving the field strength, the authors consider the problem of
finding states of the electromagnetic field in which the electrical
or the magnetic field strength has a well defined value. The states
of interest are those in which the field strength is given in all of
space at one particular time, so that a complete set of the values
of the electrical (or magnetic) field strength can be chosen in all

skoy SSR (Institute of Radiophysics and Electronics, Academy of Sciences of the Ukrainian SSR)

NR REF SOV: 003

OTHER: 002

ZAKHAROV, A.P.; VECHER, N.A.; LEKONISEV, A.N.; RUDNITSKIY, P.M.;
BOBCHALENKO, L.N.; TSHEBNIK, Z.G.; ARYASOV, N.I., inzh.,
retiree; DOVCOFOL, V.I., red.; DUBROV, I.P., red.;
GETLING, Yu., red.

[Vanadium of the Kachkanar deposit] Kachkanarskii vanadii.
Sverdlovsk, Sredne-Ural'skoe knizhnoe izd-vo, 1964. 302 p.
(MIRA 18:11)

BURDAKOV, Dmitriy Dmitriyevich; TSUKERNIK, Zinoviy Grigor'yevich; YEL'KIN, S.A., inzh., retsenzent; ROMANOV, A.A., kand. tekhn. nauk, retsenzent; BENYAKOVSKIY, M.A., inzh., retsenzent; GOL'DSRTEYN, M.I., kand. tekhn. nauk, retsenzent; DUBROV, N.F., nauchnyy red.; SYRCHINA, M.M., red. izd-va; KRYZHOVA, M.L., red. izd-va; TURKINA, Ye.D., tekhn. red.

[Metallurgy of ferrrous metals; manual for the training of skilled workers in industry] Metallurgiiia chernykh metallov; uchebnoe posobie dlia podgotovki kvalifitsirovannykh rabochikh na proizvodstv. Sverdlovsk, Gos. nauchno-tekhn. izd-vo lit-ry po chernoi i tsvetnoi metallurgii, 1961. 437 p. (MIRA 14:11)
(Iron—Metallurgy) (Steel—Metallurgy) (Metalwork)

IGNATIK, A.F., inzh.; TSUKERNIK, Z.S., inzh.

Using radioactive pickups in the automatization and regulation
of stone-materials production. Stroi.i dor.mashinostr. 4 no.9:
28-30 S '59. (MIRA 12:11)

(Radioisotopes--Industrial applications)

(Automatic control)

(Building materials industry)

IGNATYUK, A.F., inzh.; TSUKERNIK, Z.S., inzh.

Using tagged atoms for determining the speed of movement of
concrete through steel pipes. Stroi. i dor. mashinostr 3 no.5:
23-24 My '58. (MIRA 11:5)
(Radioactive tracers--Industrial applications)
(Concrete--Transportation)

TSUKERNIK, Z.S.

TSUKERNIK, Z.S., inzhener.

New tensiometering instrument. Sbor.trud.VNIISTroidorrazh
Lenfil. no.1:91-95 '57. (PIRA 10:8)
(Electronic instruments)
(Tensiometers)

PANFILOVA, T.A.; SUVOROVSKAIA, N.A.; TSUKERSHTEYN, O.E.

The coronary vasoconstrictor activity of the serum of rabbits on a cholesterol diet. Cor Vasa 6 no.4:308-311 '64.

1. The Central Research Laboratory, 1st Leningrad Medical Institute, Leningrad, USSR.

PANFILOVA, T.A.; TSUKERSHTEYN, O.Ya.

Modified spreading rate of the pulse wave in some arrhythmias under experimental conditions. Biul. eksp. biol. i med. 58 no. 7: 30-32 J1 '64. (MIRA 18:2)

1. Tsentral'naya nauchno-issledovatel'skaya laboratoriya (zav. - dotsent V.M. Ivanov) I Leningradskogo meditsinskogo instituta imeni Pavlova. Submitted June 15, 1963.

TSUKERSHTEYN, O.Ye.

Effect of vagotropic substances on certain functions of the
liver, effect of the nervous system on the liver. Ter. arkh.,
Moskva 23 no. 6:43-55 Nov-Dec. 1951. (CML 21:3)

1. Of the Naval Medical Academy.

TSUKERSHTEYN, O.Ye.; KUPSIK, B.M.

Some variants in the hemopoietic system in diphyllbothriasis.
Probl.gemat. i perel.krovi 4 no.7:49-52 J1 '59. (MIRA 12:10)

1. Iz I Leningradskogo meditsinskogo instituta imeni akad.
I.P.Pavlova.

(TAPEWORM INFECTION, compl.
diphyllbothriasis causing anemia (Rus))
(ANEMIA, etiol. & pathogen.
diphyllbothriasis (Rus))

TSUKERSHTEYN, O. YE.
TSUKERSHTEYN, O. Ye.; KUPSIK, B.M.

Some peculiarities in the composition of the blood in diphyllor-
bothriasis in the Far North. Sov.med. 21 Supplement:7 '57.
(ARTIC REGIONS--TAPEWORMS) (MIRA 11:2)
(BLOOD--EXAMINATION)

LINDENBRATEN, L.D. (Leningrad); TSUKERSHTEYN, O.Ye. (Leningrad).

Diagnosis of internal biliary fistulas; symptom of gas in bile ducts. Klin.
med. 31 no.8:38-42 Ag '53. (MLBA 6:11)

(Biliary tract) (Fistula)

TSUKERSHTEYN, O.Ye., kand.med.nauk

Case of diabetes mellitus developing under the influence of ACTH
and cortisone. Probl.endok.i gorm. no.1:115-116 '62.

(MIRA 15:8)

(ACTH)

(CORTISONE)

(DIABETES)

PANFILOVA, T.A.; TSUKERSHTEYN, O.Ye.; VOL'FE, M.M.

Correlations between functional and morphological changes in experimental cholesterol atherosclerosis in dogs. Pat. fiziol. i eksp. terap. 9 no.5:53-57 S-0 '65. (MIRA 19:1)

1. "Sentral'naya nauchno-issledovatel'skaya laboratoriya (zav. - dozent V.M. Ivanov) I Leningradskogo meditsinskogo instituta imeni I.P. Pavlova. Submitted May 23, 1964.

TSUKERSHTEYN, O. YE.

"Leukemoid Reactions During Intoxication With Tetraethyl Lead",
Military-Medical Journal, No. 8, P 34, Aug 1955.

MASLENNIKOV, V.D; TSUKERSHTEYN, O.Ye.

Leucocytosis and absolute monocytosis following Valdman's cup
endothelial test. Lab. delo 7 no.3:22-25 Mr '61.

(MIRA 14:3)

(LEUCOCYTOSIS)

(RHEUMATIC FEVER)

TSUKERSHTEYN, V.G., and RUDERMAN, A.I.

"Experimental Studies on the Distribution of X-ray Radiation Doses when Sectorial Irradiation by Means of Swinging X-ray Tubes is being applied", p. 165. Research Into the Field of X-Ray Technique, Vol. 9, Medgiz, Moscow 1955

RUDERMAN, A.I., doktor med. nauk; TSUKERSHTEYN, V.G.

Indications for X-ray treatment of esophageal cancer by sector
irradiation with a pivoting tube. Trudy TSentr. nauch.-issl. inst.
rentg. 1 rad. 10:226-230 '59. (MIRA 12:9)
(ESOPHAGUS--CANCER) (X RAYS--THERAPEUTIC USE)

TSUKERSHTEYN, V.G.

X-ray treatment of benign vascular tumors. Trudy TSentr. nauch.-
issl. inst. rentg. i rad. 10:273-278 '59. (MIRA 10:9)
(X RAYS---THERAPEUTIC USE) (TUMORS)

RUDERMAN, A.I.: TSUKERSHTEYN, V.G.

Experimental study of the distribution of X-ray emission in the case
of sectional exposure by oscillating tubes. Trudy TSentr. nauchn.
issl.inst.rentg. 1 rad. 9:165-173 '55. (MLBA 9:12)
(X RAYS)

TSUKERSHTEIN, YE. I.

Diabetes 2. izd-perer. i dop. Leningrad, Gos. ordena Lenina in-t
usovershenstvovaniia vrachei im. S.M. Kirova, 1947. 336 P.

DAFM

L. Diabetes

PA 07/49100

TSUKERSHTEYN, Ye.I. (Prof)

USSR/Medicine - Blood, Cholesterol * Mar 49
Blood, Ketone Bodies

"Presence of Ketone Bodies and Cholesterol in the Blood of Hypertonia Cases," Prof Ye. I. Tsukershteyn, Chief, Third Chair of Internal Diseases, M. N. Yegorova, Third Chair of Internal Diseases, State Ord of Lenin Inst For Advancement of Doctors (Imeni S. M. Kirov), 6 pp

"Klin Med" No 3

Gives results of studying hypertonic cases in three tables. From data obtained, it may be

65/49168

USSR/Medicine - Blood, Cholesterol Mar 49
(Contd)

assumed that hyperketonemia has a serious effect on the general condition of the patient. Changes in general or alimentary functions affect the content of ketone bodies. In most cases under proper treatment, the cholesterol content in the blood tends to approach normal as general condition and liver functions improve. Dir, State Ord of Lenin Inst for Advancement of Doctors: G. A. Znamenskiy.

65/49168

TSUKERSHTEYN, Ye.I.; YEGOROVA, M.N.

Barrier liver function in hypertension. Klin. med., Moskva 30 no.
11:90 Nov 1952. (GLML 23:5)

1. Professor for Tsukershteyn. 2. Of the Third Department of Internal Diseases (Head -- Prof. Ye. I. Tsukershteyn), State Order of Lenin Institute for the Advanced Training of Physicians imeni S. M. Kirov.

TSUKERSHTEYN, Ye.I., professor (Leningrad)

Peculiarities of the course of rheumatism in adolescents. *Klin. med.*
31 no.11:48-52 N '53.

(MLRA 6:12)
(Rheumatism)

TSUKERSHTEYN, Ye.I., prof.; TNIMOVA, K.Z., dotsent

Peculiarities in the clinical aspects and treatment of tuberculosis of the lymph nodes of the abdominal cavity. Zdrav. Kazakh. 21 no.1: 43-48 '61. (MIRA 14:3)

1. Iz kafedry gosspital'noy terapii (zav. prof. Ye.I. Tsukershteyn) Karagandinskogo meditsinskogo instituta. (LYMPHATICS—TUBERCULOSIS) (ABDOMEN)

TSUKERSHTEYN, Ye.I., prof.; KARLINSKIY, V.M., kand.med.nauk

Prevention of rheumatic fever in adults. *Zdrav. Kazakh*, 21 no.5:
5-8 '61. (MIRA 15:2)

1. Iz kafedry gospital'noy terapii Karagandinskogo meditsinskogo
instituta.

(RHEUMATIC FEVER)

PROCESSES AND PROPERTIES INDEX

CL-4

BC

Composition of (A) *Ferula drupacea*, L., and of the oil pressed from its seeds, (B) roots of *Ferula pseudoreconitana*, Lipsky. 1. TRUMAN, YANIK and V. BARSUKAI (Bull. Univ. Aste Centrale, 1933, No. 21, 49-54, 55-66).—(A) The air-dry leaves contain hemicellulose (I) 7.7, cellulose (II) 9.3, lignin (III) 5.5, protein 16.8, ash 8.7, H₂O-sol. 24, Et₂O-sol. substances 13.8%, the roots contain 8.9% of H₂O-sol. products, and the seeds contain (I) 6.8, (II) 8.3, (III) 4.1, ash 6.0, H₂O-sol. 23.5, EtOH-sol. 2.1, and Et₂O-sol. substances 15.4%. The Et₂O extract of the seeds yields a fatty oil, containing saturated (palmitic, stearic) 8, and unsaturated acids (oleic, linoleic, linolenic) 80, resins 4.1, and unsaponifiable 5.0%.

(B) [with C. V. BRUTKVA and B. I. AJEKOVITSCII]. The air-dry root contains (I) 11, (II) 12.4, (III) 4.3, pentosans 6.4, pectins 9.6, starch 7.19, mono- 2.11, and di-saccharides 8.89, protein 7.37, ash 5.82, H₂O-sol. 29.8, EtOH-sol. 1.9, and Et₂O-sol. substances 15.9% (including resin 12.63 (acids, including podocarpin, 6.1, phenols 62.5, and neutral constituents 29.5%), essential oil 3.3 (*d*-α-pinene 90, *d*-β-phellandrene 5%)).

R. T.

ASS-SLA METALLURGICAL LITERATURE CLASSIFICATION

FROM SOURCE

FROM SOURCE

1ST AND 2ND ORDERS PROCESSES AND PROPERTIES INDEX 3RD AND 4TH ORDERS

Ca

13

Utilization of cottonseed husks for synthesis of glucose, lactic acid and glycerol. I. P. Tsukerivanik, P. P. Samsonov, L. P. Danilevskii and N. G. Sidorova. *Bull. univ. Asie centrale* No. 31, 60-82(1935).—One kg. of husks is heated for 1 hr. at 20° with 1 l. of 75.5% H₂SO₄, undissolved material is removed, 8 l. of H₂O added, and the soln. is boiled (for 3 hrs.) and made neutral with CaO or CaCO₃. Glucose is obtained in 27-30% yield from the condensed filtrate and may be fermented to yield lactic acid, or EtOH and glycerol. B. C. A.

ASB-51A METALLURGICAL LITERATURE CLASSIFICATION

MATERIALS INDEX COMMON ELEMENTS COMMON LITERATURE INDEX

GROUPS	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
--------	---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----

10

Alkylation of phenols with alcohols in the presence of aluminum chloride. II. Alkylation with secondary and primary aliphatic alcohols. I. P. Tsukerivanik and Z. N. Nazarova. *J. Gen. Chem.* (U.S.S.R.) 7, 1023-24 (1937); *C. A.* 30, 4431. The difficulties of obtaining alkylphenols by catalytic condensation of phenols with secondary alcs. have been overcome by the use of excess $AlCl_3$ and higher temps. By this method about 70% alkylphenols, contg. small amts. of dialkylphenol ethers (II), $(R_2C_6H_3OR)$ (I) and alkylphenol ethers (RC_6H_4OR) (II), are formed. The proportion of II in the reaction mixt. increases with decrease in the $AlCl_3$ concn. and temp. The probable mechanism of the condensation is: $ROH + AlCl_3 \rightarrow HCl + AlCl_2(OR)$; $PhOH + AlCl_2(OR) \rightarrow HCl + PhOAlCl_2$; $AlCl_2(OR) + PhOAlCl_2 \rightarrow PhOR + AlCl_3 + AlCl_2OH$. PhOR is then converted to II either by alkylation: $PhOR + ROH + AlCl_3 \rightarrow II + AlCl_2OH + HCl$, or by rearrangement: $2 PhOR \xrightarrow{AlCl_3} II + PhOH$ (cf. Smith, *C. A.* 27, 5067). The alkylated ethers are then converted to alkylphenols by the action of the excess of $AlCl_3$ or higher temps. either by sapon.: $II + HCl \xrightarrow{AlCl_3} RC_6H_4OH + RCl$ (Hartmann and Gattermann, *Ber.* 25, 3531 (1892)), or by rearrangement: $II + PhOH \xrightarrow{AlCl_3}$

$2RC_6H_4OH + 2 H \xrightarrow{AlCl_3} RC_6H_4OH + H$, (cf. Smith, *loc. cit.*). Condensation of PhOH with $MePhCHOH$ resulted chiefly in a mixt. of *o*- and *p*- $MePhCHOH$ and a little of $Et_2C_6H_3OH$ as a result of the isomerization of the alk. radical, which could be effected either by the cleavage of H_2O from the alk. by the action of $AlCl_3$ or by the isomerization of alkylated phenol ether. The condensation of PhOH with primary alcs. was effected by the same method with 2 mols. of $AlCl_3$ to 1 mol. of PhOH and 1 mol. of alc. Condensation of Me_2CHOH with PhOH in petr. ether at $110-20^\circ$ for 6 hrs. gave 52% *p*- $HOC_6H_4CHMe_2$ (III), m. 61° , (the ester m. $71-2^\circ$), and about 23% of nonphenolic fraction, b. $225-30^\circ$, contg. chiefly *p*- $Me_2CHOC_6H_4CHMe_2$ (IV). Anisole and Me_2CHOH in petr. ether at 120° for 4 hrs. gave 38% of a mixt. of III and *o*-isost. er, b. $215-17^\circ$, and 50% of *o*- and *p*- $MeOC_6H_4CHMe_2$ contaminated by a little IV. The condensation at $130-40^\circ$ for 5-6 hrs. resulted in 64% of *o*- and *p*- $HOC_6H_4CHMe_2$ and 30% $MeOC_6H_4CHMe_2$. *m*-Cresol and Me_2CHOH heated at 125° for 6 hrs. gave 68% *3,4-HOC_6H_3Me_2CHMe_2, m. 112° , and 20% *3,4-Me_2CHOC_6H_3Me_2CHMe_2*, b. $233-5^\circ$. PhOH and $EtMeCHOH$ in ligrom*

see other side \rightarrow

ASS-51A METALLURGICAL LITERATURE CLASSIFICATION

heated at 120-40° for 6 hrs. gave 52% of EtMeCHC₆H₄OH
o-isomer, b. 229-32°, d₄ 0.9007, n_D 1.5182, M. R. 45.89;
p-isomer, b. 235-7°, d₄ 0.8993, n_D 1.518, and 13% Et-
 MeCHOC₆H₄CHMeEt. In the process of distn. the
 ether is given into the *tert*-butylphenol. Anisole and
 EtMeCHOH digested on a water bath for 2 hrs. gave 83%

MeOC₆H₄CHMeEt, b. 109-70°, d₄ 0.8413, n_D 1.503,
 M. R. 61.5, 16% MeOC₆H₄(CHMeEt)₂, b. 140-2°, d₄
 0.9240, n_D 1.498, M. R. 99.72, and 13% HOOC₆H₄CHMeEt.
 PhOH and MePrCHOH allowed to stand at room temp.
 for 24 hrs. and then heated at 40-50° for 4 hrs. gave 57.5%
 of phenolic and 11% of nonphenolic fraction. From the
 phenolic fraction crystd. out Et₂CHC₆H₄OH, m. 88° (Bz
 deriv., m. 40-1°); the residue on fractionation gave *p*-
 MePrCHC₆H₄OH, b. 150-6° (Bz deriv., b. 255°), and
o-isomer, b. 140-50°, d₄ 0.8981, n_D 1.519 (Bz deriv., b.
 340-4°). The nonphenolic fraction gave MePrCHC₆H₄-
 OCHPrMe, b. 230-40°, and (MePrCH)₂C₆H₄OCHPrMe,
 b. 245-60°, d₄ 0.900, n_D 1.4919. Anisole and MePr-
 CHOH treated as above afforded MeOC₆H₄CHPrMe, b.
 221-50°, d₄ 0.9284, n_D 1.499, and MeOC₆H₄(CHPrMe)₂,
 b. 245-60°. The condensation of PhOH and also EtOH heated at

120-40° for 6 hrs. gave 62% phenolic and 11% nonphenolic
 products. The phenolic fraction consisted of 60% Et-
 C₆H₄OH, b. 237°, m. 76-8.5°, and 40% *o*-EtC₆H₄OH,
 m. 137°, and the *p*-isomer, b. 216-20°. The nonphenolic
 fraction on redistn. was completely converted into Et₂C₆-
 H₄OH. PhOH and PrOH gave 73.6% *o*-PrC₆H₄OH, b.
 221°, d₄ 1.0006, and the *p*-isomer, b. 225-8°, and 17%
 PrC₆H₄OPr, b. 340-5°; this on standing was converted
 into PrC₆H₄OH. PhOH and BuOH gave 72% phenolic
 and 14% nonphenolic products. *o*-BuC₆H₄OH, b. 233-5°,
 d₄ 0.9713; *p*-isomer, b. 240-60°, d₄ 0.9761 (Bz deriv.,
 m. 25-6°). PhOH and *tert*-BuOH gave C₆H₅ and 82%
 Me₂CC₆H₄OH, m. 98°. PhOH and *iso*-AmOH gave 3
 amylphenols, of which only Me₂RtCC₆H₄OH, m. 28-90°,
 was identified. About 25 references. Chas. Blanc

(CONTINUED) PROCEEDINGS

GA Condensation of alcohols with aromatic hydrocarbons in the presence of anhydrous aluminum chloride. III. Condensation of primary alcohols with benzene and toluene. I.P. Tsukervanik and G. Vikhrova. J. Gen. Chem. (U.S.S.R.)

10

In the presence of anhydrous AlCl3 and PhMe, the condensation of primary alcohols with benzene and toluene proceeds by intermediate cleavage of H2O with the formation of cycloalkenes and cycloalkyl chlorides: C6H5OH + AlCl3 -> C6H5 + AlCl3OH + HCl; C6H5 + HCl -> C6H5Cl

the condensation of secondary cyclic alcs. with aromatic hydrocarbons proceeds by intermediate cleavage of H2O with the formation of cycloalkenes and cycloalkyl chlorides: C6H5OH + AlCl3 -> C6H5 + AlCl3OH + HCl; C6H5 + HCl -> C6H5Cl

isomer (identified as m-C6H4(CO2Me)2, m. 62°) and sym-C6H2(C6H11)3, m. 68°, identified as tri-Me trimesate, m. 144°. PhMe is less reactive, requiring more AlCl3 and a large excess of PhMe. C6H5OH (70 g.), 80 g. AlCl3 and 200 cc. PhMe gave 72% of a fraction, bp 254-60°, d420 0.9343, nD20 1.5242, M. R. 57.95, consisting of p- and m-MeC6H4C6H11, identified as p- and m-C6H4(CO2Me)2, resp., and 18% 3,5-Me2C6H3(C6H11), m. 93.5°, identified as tri-Me trimesate. From the decomps. of C6H5OH with AlCl3 with the formation of 16-20% C6H5 and 20-5% C6H5Cl it is assumed that by analogy with tertiary alcs.,

ASB-31A METALLURGICAL LITERATURE CLASSIFICATION

Table with columns for document number, date, and classification codes. Includes fields like '440000 #', '101000 WIP Date Date', and '43110001'.

10

PROCESSES AND PROPERTIES INDEX

ca

Condensation of alcohols with aromatic hydrocarbons in the presence of anhydrous aluminum chloride. VI. Alkylation of halogenated aromatic hydrocarbons by alcohols. I. P. Tsukervanik. *J. Gen. Chem.* (U. S. S. R.) 8, 1512-15 (1939); cf. *C. A.* 31, 5789. —The preliminary exptl. data on the condensation of excess of PhCl, *o*-ClC₆H₄Me (I) and 1-C₆H₄Cl (II) with alcs. in the presence of AlCl₃ at elevated temps. show that the alkylation of aryl halides proceeds analogously to that of unsubstituted aromatic hydrocarbons with the formation of corresponding monoalkyl- (47-65% yield) and polyalkylaryl chlorides. The alkyl derivs. represent mixts. of *p*- and some *m*-isomers, the constitution of which was detd. by oxidation to the corresponding chlorobenzoic acids. The condensation with primary alcs. (EtOH) proceeds most difficultly (requiring a large excess of AlCl₃ (1.5 g./mol.); with secondary alcs. less difficultly (1 g./mol. AlCl₃); and with tertiary alcs. very easily (0.5 g./mol. AlCl₃). Greater AlCl₃ concn. and higher temps. increase the formation of polyalkyl derivs. The reactions are effected by gradual addns. of AlCl₃ to a mixt. of an aryl chloride and an alc. and refluxing on a boiling water bath for 2-3 hrs. Unless otherwise stated, the values given below are for *b_{mp}* and *d₄²⁰*, resp. The reaction of 37 g. *tert*-BuOH and 100 g. PhCl with 30 g. AlCl₃ gave 11 g. polyalkyl chlorides (III) and 65% of isomeric *tert*-BuC₆H₄Cl; *p*-isomer, 210-12°, *n_D²⁰* 1.5125. PhCl (100 g.) and 44 g. *tert*-AmOH with 25 g. AlCl₃ gave 8 g. III and 50% *p*- and *m*-*tert*-AmC₆H₄Cl; *p*-isomer, 224-9°, *n_D²⁰* 1.5139. PhCl (100 g.) and 60 g. iso-PrOH with 120 g. AlCl₃ gave 62% chlorocumenes; *p*-isomer, 195-7°, *n_D²⁰* 1.514. PhCl (100 g.) and 44 g. *sec*-BuOH with 30 g. AlCl₃ gave 8 g. III and 60% isomeric *sec*-BuC₆H₄Cl; *p*-isomer, 212-14°, *n_D²⁰* 1.4995. PhCl (100 g.) and 30 ml. EtOH with 120 g. AlCl₃ gave 10 g. III and 40% *p*-EtC₆H₄Cl, 179-80°, *n_D²⁰* 1.5235. PhCl (150 g.) and 74 g. iso-BuOH with 200 g. AlCl₃ gave 25 g. III and 30% of monoalkyl derivs., from which was isolated *p*-*tert*-BuC₆H₄Cl. PhCl (100 g.) and 44 g. iso-AmOH with 80 g. AlCl₃ gave 15 g. III and 35% monoalkyl derivs.; *p*-*tert*-AmC₆H₄Cl, 224-30°, *n_D²⁰* 1.5139. I (25 g.) and 19 g. *tert*-BuOH with 15 g. AlCl₃ gave 45% monoalkyl derivs.; *tert*-BuMeC₆H₄Cl, 225-30°, *n_D²⁰* 1.5145. II (120 g.) and 44 g. *tert*-AmOH with 30 g. AlCl₃ gave 15 g. III and 60% monoalkyl derivs.; *tert*-amyl-1-chloronaphthalene, 310-12°, *n_D²⁰* 1.1235, *n_D²⁰* 1.0099. II (75 g.) and 30 g. iso-PrOH with 60 g. AlCl₃ gave 12 g. III and 45% iso-propyl-1-chloronaphthalene, *b. 305-10°*, *n_D²⁰* 1.1421, *n_D²⁰* 1.0043. Chas. Blanc

ASB-51A METALLURGICAL LITERATURE CLASSIFICATION

123000 1157 0001 001

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	00
---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----

TSUKERVANIK, I. P.

Benzylstibine and its derivatives. I. P. Tsukervanik and D. Sukhray. *J. Gen. Chem.* (U. S. S. R.) 7, 1527-31 (1937). — PhCH_2MgCl and SnCl_4 in a CO_2 atm. give *tri-benzylstibine* (I), m. 90° , which is easily oxidized by air, especially in soln., to *tribenzylstibine oxide* (II). I reacts with Cl_2 and Br_2 , or II with HCl and HBr , to give the *dichloride* m. 100° , and the *dibromide* m. 102° , resp. When I is heated with SnCl_4 , it forms $(\text{PhCH}_2)_3\text{SnCl}$, m. 187° , which gives with dil. Na_2CO_3 soln. $[(\text{PhCH}_2)_3\text{SnO}]_2$. With HBr this forms $(\text{PhCH}_2)_3\text{SnBr}$, m. 150° (decomp.). H. M. Leicester

16

Condensation of alcohols with aromatic compounds in the presence of aluminum chloride. IX. Condensation of 1,2-methylcyclohexanol, menthol and borneol with benzene and toluene. I. P. Tsakeryanik and N. G. Sidonova, *Gen. Chem. (U. S. S. R.)* **8**, 1800 (1962) (in English, 1962) (1968); cf. *C. A.* **33**, 4587¹. The following compounds were prepd.: *methylcyclohexylbenzene*, $C_{11}H_{18}$ (42.8% yield), b_p 142.3°, d_4^{20} 0.8608, n_D^{20} 1.5290; *(methylcyclohexyl)toluene* isomers (50% yield), b_p 152.4°, d_4^{20} 0.8590, n_D^{20} 1.5205; *menthylbenzene* $C_{11}H_{18}$ (1) (51.5% yield), b_p 151.4°, d_4^{20} 0.8642, n_D^{20} 1.5207; *p*- and *m*-*menthyltoluene* (61% yield), b_p 183.8°, d_4^{20} 0.8244, n_D^{20} 1.5185; *bornylbenzene*, $C_{11}H_{18}$ (30% yield), b_p 165.8°, d_4^{20} 0.8535, n_D^{20} 1.5250; *p*- and *m*-*bornyltoluene* (64% yield), b_p 131.3°, d_4^{20} 0.8500, n_D^{20} 1.5275. In the prepn. of 1,3-menthene, b_p 165.0°, d_4^{20} 0.8103, n_D^{20} 1.4170, and *tert*-menthylchloride, $C_{11}H_{17}Cl$, b_p 117.8°, d_4^{20} 0.8105, n_D^{20} 1.4070, were the chief products when insufficient $AlCl_3$ was used. Small amts. of polyalkylated hydrocarbons were always obtained but not identified. John Livak

ASB, S.E.A. METALLURGICAL LITERATURE CLASSIFICATION

830M SYMBLIVM

NO.	SYMBOL	CLASSIFICATION	DATE	STATUS
1	11	11	11	11
2	11	11	11	11
3	11	11	11	11
4	11	11	11	11
5	11	11	11	11
6	11	11	11	11
7	11	11	11	11
8	11	11	11	11
9	11	11	11	11
10	11	11	11	11
11	11	11	11	11
12	11	11	11	11
13	11	11	11	11
14	11	11	11	11
15	11	11	11	11
16	11	11	11	11
17	11	11	11	11
18	11	11	11	11
19	11	11	11	11
20	11	11	11	11
21	11	11	11	11
22	11	11	11	11
23	11	11	11	11
24	11	11	11	11
25	11	11	11	11
26	11	11	11	11
27	11	11	11	11
28	11	11	11	11
29	11	11	11	11
30	11	11	11	11
31	11	11	11	11
32	11	11	11	11
33	11	11	11	11
34	11	11	11	11
35	11	11	11	11
36	11	11	11	11
37	11	11	11	11
38	11	11	11	11
39	11	11	11	11
40	11	11	11	11
41	11	11	11	11
42	11	11	11	11
43	11	11	11	11
44	11	11	11	11
45	11	11	11	11
46	11	11	11	11
47	11	11	11	11
48	11	11	11	11
49	11	11	11	11
50	11	11	11	11
51	11	11	11	11
52	11	11	11	11
53	11	11	11	11
54	11	11	11	11
55	11	11	11	11
56	11	11	11	11
57	11	11	11	11
58	11	11	11	11
59	11	11	11	11
60	11	11	11	11
61	11	11	11	11
62	11	11	11	11
63	11	11	11	11
64	11	11	11	11
65	11	11	11	11
66	11	11	11	11
67	11	11	11	11
68	11	11	11	11
69	11	11	11	11
70	11	11	11	11
71	11	11	11	11
72	11	11	11	11
73	11	11	11	11
74	11	11	11	11
75	11	11	11	11
76	11	11	11	11
77	11	11	11	11
78	11	11	11	11
79	11	11	11	11
80	11	11	11	11
81	11	11	11	11
82	11	11	11	11
83	11	11	11	11
84	11	11	11	11
85	11	11	11	11
86	11	11	11	11
87	11	11	11	11
88	11	11	11	11
89	11	11	11	11
90	11	11	11	11
91	11	11	11	11
92	11	11	11	11
93	11	11	11	11
94	11	11	11	11
95	11	11	11	11
96	11	11	11	11
97	11	11	11	11
98	11	11	11	11
99	11	11	11	11
100	11	11	11	11

10

CA

PROCESSES AND PROPERTIES INDEX

Reciprocal action of acetylketene and alcohols. I.
 I. P. Tsukervanik and I. A. Ermolenko. *Bull. univ. Acad. Sci. Ser. Chem. Div.* 22: 215-20 (in German, 220) (1938).—The reaction between acetylketene and various alcs. and some other compts. has been investigated. The acetylation proceeds even in the cold in the absence of catalysts. The catalysts tested were quinoline, H₂SO₄, and ZnCl₂. Only in acetylating glycerol the presence of H₂SO₄ is of advantage. The reaction proceeds smoothly and the yield is almost theoretical in the case of primary and secondary alcs., whereas the acetylation of tertiary alcs. proceeds much more slowly. Glycerol and glycol give a mixt. of mono- and diacetates. Unsatd. alcs. react only with the HO group with the formation of normal acetates. The acetylation of *p*-anisidine gives 98% of pure acetate. Aspirin is obtained in 96% of the theoretical yield from salicylic acid.
 Gertrude Berend

METALLURGICAL LITERATURE CLASSIFICATION

ASB-31A

10

C.F.

Alkylation of phenol and anisole by the Friedel-Crafts method. I. P. Tsukerman and V. D. Fomichov. *Bull. univ. Mosc. chim. ser.* 22, 221-4 (in German, 225) (1938). - Alkylation of phenol (I) and anisole (II) has been carried out with BuCl (III), iso-BuCl (IV) and iso-AmCl (V) in the presence of an equimol. amt. of AlCl₃. In the case of II, the most favorable conditions were 4 hrs. heating of the mixt. on the water bath, using ligroin as a solvent, whereby a mixt. of *o*- and *p*-alkylanisole is obtained in 70% of the theoretical yield and small amts. of dialkylanisoles. Sapon. of II occurs simultaneously to some extent with the formation of phenol. In the case of I, longer heating to higher temps. is required. The best yield, up to 60% of the theory, is obtained in the absence of a solvent. In addn. to a mixt. of *o*- and *p*-alkylphenols, alkyl ethers of phenol are formed. To 25 g. II, 20 g. V and 100 ml. ligroin, 35 g. AlCl₃ is added slowly within 1 hr. and the mixt. heated for 4 hrs. on a boiling water bath. Isoamylanisole (10 g.), b_m 128-2°, d₄²⁰ 0.9401, n_D²⁰ 1.4995, and 3 g. diisoamylanisole, b_m 137-40°, d₄²⁰ 0.9511, n_D²⁰ 1.5112, are formed. II (22 g.), 20 g. III, 35 g. AlCl₃, and 100 ml. ligroin when heated for 4 hrs. on a boiling water bath give 21 g. butylanisole from which is isolated *p*-butylanisole, b_m 108-10°, d₄²⁰ 0.9273, n_D²⁰ 1.5020, and 3 g. dibutylanisole, b_m 130-7°, d₄²⁰ 0.9288, n_D²⁰ 1.5105. By heating 60 ml. II with 20 g. IV and 27 g. AlCl₃ there is formed 13 g. isobutylanisole, b_m 126-7°, d₄²⁰ 0.9572, n_D²⁰ 1.4980. I (19 g.), 20 g. III and 27 g. AlCl₃ on heating for 4 hrs. to 110° give 9 g. butylphenol, from which is isolated the *p*-isomer, b_m 129-30°, d₄²⁰ 0.9749, n_D²⁰ 1.5180. 5 g. I and 4 g. of butylphenol Bu ether, b_m 144-7°, d₄²⁰ 0.9049, n_D²⁰ 1.5000. I (19 g.), 22 g. V and 27 g. AlCl₃ heated for 3 hrs. on a boiling water bath give 4 g. isoamylphenol, b_m 130-3°, n_D²⁰ 1.5202, 10 g. I and 6 g. of isoamylphenol isoamyl ether. When AlCl₃ is added with cooling, 55% *tert*-amylphenol, m. 92°, are obtained instead of isoamylphenol. Gertrude Berend

Condensation of aldehydes and ketones with aromatic hydrocarbons in the presence of anhydrous aluminum chloride. I. Condensation of aliphatic ketones with phenols. I. P. Tarkervanik and Z. N. Nazarova. *J. Gen. Chem. (U. S. S. R.)* 9, 83-5(1939); cf. *C. A.* 31, 5778. The interaction of aliphatic ketones with PhOH in the presence of excess AlCl₃ (1.5 mols.) in the cold for 1-4 days yields dialkyl diphenolmethanes, which on heating decomp. to give *ortho*-alkyl phenols. The latter are formed directly in 40% yield by condensation of the reaction mixt. on the water bath for 3-4 hrs. The mechanism of the condensation reaction is analogous to the similar procedure in the presence of fuming HCl studied by Diamm (*J. Russ. Phys.-Chem. Soc.* 23, 488, 523, 601(1891); cf. *loc. cit.*). In this way were prepd. *isopropylphenol* (from Me₂CO and PhOH), b. 200-30° (and considerable resinification products) and *sec-amylphenol* (I) (from Et₂CO), m. 79°, b. 245-80° (*Be deriv.*, m. 48-50°). Me₂COPr also formed I, b. 245-50°, d₄²⁰ 0.97, n_D²⁰ 1.5215. It gives a *Be deriv.*, b. 340-50°, *Ac deriv.*, b. 251-5°, and *Me ether*, b. 232-8°, which with Cl₂ is oxidized to anisic acid, m. 170°. Condensation of *m*-cresol with Me₂CO produced an anhydride, m. 131°, described by Zincke and Gaebel (*C. A.* 6, 1753). Chas. Blanc

ASD S.A. METALLURGICAL LITERATURE CLASSIFICATION

COMMON ELEMENTS

10

Alkylation of aromatic hydrocarbons by dihalogen derivatives. I. Condensation of 1,3-chlorobromopropane with benzene. I. Tsukervanik and K. Yatsimirskii. *J. Gen. Chem. (U. S. S. R.)* 10, 1075 (1940). When $\text{Cl}(\text{CH}_2)_2\text{Br}$ and C_6H_6 react in the presence of AlCl_3 at $0-12^\circ$, 40% $\text{Ph}(\text{CH}_2)_2\text{Br}$ (I) is formed. At $80-5^\circ$, 60% $\text{Ph}(\text{CH}_2)_2\text{Ph}$ is formed, with some PhPh as a by-product, but no I is obtained. H. M. Leicester.

Lab. of Organic Chem., Cen. Resin State Univ.

ASSOCIATED METALLURGICAL LITERATURE CLASSIFICATION

WATERGATE INDEX

LIST AND THE GROUPS RECEIVES AND PROPERTIES INDEX 100 AND 4TH CROSS

ca

10

The alkylation of aromatic compounds by alcohols in the presence of anhydrous ferric chloride. Z. N. Nazarov and I. P. Tsukeryank, *J. Gen. Chem. (U. S. S. R.)* 10, 1181-5(1940).—Tertiary alcs. react easily at room temp. with C_6H_5 , $MePh$, C_6H_5 , $PhOH$, *m*- and *p*-cresols and $MeOPh$ to give 70-85% yields of monoalkyl derivs. in the presence of 0.5 equiv. $FeCl_3$ per equiv. of alc. If the mixt. is heated, some *p*-alkyl compds. are also formed. With secondary alcs., a 1:1 mixt. of $FeCl_3$ and alc. is required and the mixt. must be heated to 60-60° to get 38-78% yields of *o*- and *p*-derivs. With primary alcs., rate of reaction depends on structure, branched-chain alcs. reacting faster than normal alcs. H. M. Leicester

A10-51A DETAILING LITERATURE CLASSIFICATION 6-27-57 (100)

GROUP	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
-------	---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----

Condensation of alcohols with aromatic hydrocarbons in the presence of aluminum chloride. Condensation of cycloheptanol with benzene and toluene. N. G. Sidova and I. E. Zhukerzhanik. *J. Gen. Chem.* (U. S. S. R.) 10, 2073-6 (1940). - A yield of 40-50% of cycloheptylbenzene (I) and cycloheptyltoluene (II) was obtained on condensation of cycloheptanol (III) with benzene (IV) and toluene (V). III was prepd. by the reduction of cycloheptanone by the method of Mosettig and Burger (C. I. 24, 4767); 10 g. of III, 12 g. AlCl₃ and 100 ml. IV yielded 7 g. I, b.p. 132.5°, d₄²⁰ 0.8410, n_D²⁰ 1.5217, M_RD 50.42, calcd. 50.43. Oxidizing I with 15% HNO₃ in sealed tubes at 140-80° yielded II(OH), m. 119°. *p*-Cycloheptylnitrobenzene (VI) was obtained on nitrating I with a nitration mixt. VI, b.p. 203-10°. Oxidizing VI with 15% HNO₃ in sealed tubes gave *p*-O₂NC₇H₁₃CO₂H, m. 200°. The *p*-amino deriv. (VII) of I was obtained on reduction of VI with Sn and HCl, and was purified by steam distn. The benzoate of VII (from alc.) m. 173°; acetate (from deriv. of phenylcyclohexylmethane (VIII) and 1-methyl-2-phenylcyclohexane (IX), which were specially prepd., differed from the derivs. of I. *Synthesis of VIII.* Phenylcyclohexylcarbinol (X) was obtained by Grignard's reaction from 10 g. Mg, 68 g. bromocyclohexane, 5.4 g. BuH in 100 ml. abs. ether. The yield was 34.5 g. (45.5%), b.p. 130-42°. Ph cyclohexyl ketone (XI) was formed by the oxidation of X. To 31.5 g. of the heated X was added dropwise 18.5 g. Na₂Cr₂O₇ in 45 ml. of H₂O and 13

ml. H₂SO₄. The reaction product was acid with ether. After the ether was evapd. thin needles of XI, m. 54°, were obtained. The yield was 31 g. (91%). Phenylcyclohexylmethane (XII) was obtained by the reduction of XI by the Clemmensen method. The yield was 52%, M_RD 137°, d₄²⁰ 0.9430, n_D²⁰ 1.5253, M_RD 50.58, calcd. 50.43. The (nitrophenyl)cyclohexyl ethane, b.p. 187° with Sn and HCl. The *N*-H₂ deriv. (from dil. alc.) m. 101-2°. The *N*-Ac deriv. (from dil. alc.) m. 170-1°. *Synthesis of IX.* 3-Methyl-2-phenylcyclohexane (XIII) was obtained by Grignard's reaction from 1 g. Mg, 20 g. bromobenzene and 18 g. of 2-methylcyclohexanone in 50 ml. of abs. ether. The yield was 12 g. (42%), b.p. 112-15°. 1-Methyl-2-phenylcyclohexane (XIV) was obtained on heating 12 g. of XIII with 12.5 ml. HI (d. 1.7) and 0.75 g. of red P in a sealed tube, for 6 hrs., at 160-200°. XIV, b.p. 91-2°, d₄²⁰ 0.9290, n_D²⁰ 1.5300, M_RD 50.75, calcd. 50.43. The nitro and the corresponding amino compounds were obtained from XIV by the usual method. The *N*-H₂ deriv., m. 100-74°. The *N*-Ac deriv. was a liquid, b.p. 195-200°. It was obtained on treating 5 g. of III with 50 ml. of V and 6 g. of AlCl₃. The yield was 4 g. of II (a mixt. of *m*- and *p*-isomers), b.p. 110-12°, d₄²⁰ 0.9350, n_D²⁰ 1.5253, M_RD 50.81, calcd. 51.05. It oxidized with dil. HNO₃ yielded a mixt. of *tert*- and isophthalic acids, which were obtained by their di-Me esters. The conclusion is drawn that the condensation of III with IV and V proceeds only to a small extent.

James J. Lichten

AS 7-51.4 METALLURGICAL LITERATURE CLASSIFICATION

ABSTRACTS AND PROPERTIES INDEX

10

ca

Alkylation of aromatic compounds with alcohols in the presence of anhydrous FeCl₃. Z. N. Nazarova and I. P. Tsuberavnik, *J. Gen. Chem. (U. S. S. R.)* 14, 77 (1944) (English summary).—Attempts to effect a condensation between benzene and PhOH on one hand and primary alcs. (EtOH, PrOH and BuOH) on the other hand failed to give more than traces of alkaryl derivs. PhCH₂OH (30 g.), benzene (50 g.) and 3.1 g. FeCl₃ heated to 70° for 4 hrs. gave 58% Ph₂CH₂, b. 200-5°, m. 27°. PhCH₂OH (20 g.), PhOH (20 g.) and 15 g. FeCl₃ mixed, let stand for 24 hrs. and heated on a steam bath for 2 hrs. gave 58% *benzylphenol*, b. 300-30°, m. 82°. Allyl alc. (1A g.), 40 g. benzene and 20 g. FeCl₃ were mixed with cooling, let stand for 24 hrs. and finally heated for 2-3 hrs. on a steam bath to 100° to yield 3% crude *allylbenzene* and 52% (*2-chloropropyl*)benzene; the purified compds. had the following consts., resp.: b. 154-6°, n_D²⁰ 1.5110, d₄²⁰ 0.8660, and brn 200-5°, n_D²⁰ 1.5147, d₄²⁰ 1.0249. This reaction conducted at room temp. (3 days' standing) gave the 2 products in reverse ratio: 33% and 8%, resp. Allylbenzene yields a *tribromide*, m. 118°. Similarly, PhMe gave *allyltoluene*, b. 178-80°, n_D²⁰ 1.5085, d₄²⁰ 0.8618, and (*2-chloropropyl*)toluene, b. 220-30°, n_D²⁰ 1.5194, d₄²⁰ 0.8654. G. M. Kosolapoff

A.S.T.M. METALLURGICAL LITERATURE CLASSIFICATION

E-277001-28200

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----

PROCESSES AND PROPERTIES INDEX

10

Ca

The mechanism of alkylation reactions under the influence of anhydrous $FeCl_3$. Z. N. Nazarova and I. P. Tsukervanik. *J. Gen. Chem.* (U.S.S.R.) 14, 236-44 (1944) (English summary); cf. *C.A.* 39, 916¹.— H_2Br and $FeCl_3$ form HBr and $EtCl$. $PrBr$ gives $iso-PrCl$, and $iso-AmBr$ gives C_4H_{10} and mixts. of $AmCl$ and $AmBr$. When $RtBr$ and C_6H_6 are heated with $FeCl_3$, HCl , $EtCl$, and $PhBr$ are formed. BtI gives an analogous reaction. Under similar conditions $PrBr$ gives $iso-PrCl$, $PhBr$, $Br-C_6H_4Pr$, and a little $PhPr$; $iso-BuBr$ gives $PhBr$, $tert-BuPh$, $iso-BuPh$ and a mixt. which probably contains BrC_6H_4Bu , and $Bu_2C_6H_4$. $BuCl$ and $AmCl$ give almost no reaction with C_6H_6 and $FeCl_3$. $EtOH$ and $FeCl_3$ give 87% $EtCl$. With $FeCl_3$, $PrOH$ gives $iso-PrCl$ and a little $PrCl$. $BuOH$ gives $BuCl$ and some $sec-BuCl$. $iso-BuOH$ gives $tert-BuCl$ and diisobutylene. $FeCl_3$ dissolves in $iso-AmOH$, and on standing very hygroscopic crystals sep. which m. 48° , evolve HCl at 58° , and solidify at 105° to a green, nonhomogeneous mass. $iso-PrOH$ gives $iso-PrCl$; $tert-BuOH$ gives C_4H_{10} , $sec-BuCl$, and $tert-BuCl$; $tert-AmOH$ gives C_4H_{10} , and $tert-AmCl$. $iso-PropH$ is isomerized by $FeCl_3$ to PrC_6H_5OH and much tar is formed. These reactions are interpreted in terms of formation of an ionized complex of the type $(ROH.FeCl_3)^+Cl^-$ which decomps. in various ways to give the products observed. Alkylation with $FeCl_3$ is thus not a simple condensation, as with $AlCl_3$, but involves rupture of C bonds and deep-seated changes in structure. H. M. Leicester

Lab. Organic Chem. University, State Lenin, -1943-

ASB-51A METALLURGICAL LITERATURE CLASSIFICATION

GROUP #	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
---------	---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----

1ST AND 2ND ORDERS 100 AND 4TH ORDERS

PROCESSED AND PROPERTIES INDEX

10

ca

Alkylation of aromatic compounds in the presence of phosphoric acid. I. Condensation of alcohols with aromatic hydrocarbons and halides. I. P. Tsukeryanik (Middle Asiatic State Univ.). *J. Gen. Chem.* (U.S.S.R.) 15, 599-703 (1945) (English summary).—It was shown that H_3PO_4 may be used for alkylation of aromatic compds. by alcs., with yields up to 80-90% being obtained in the best cases. Considerable poly-substitution occurs, *p*-isomers predominating in disubstituted products, with occasional traces of the *o*-isomers. The iso alcs. usually isomerize into tertiary radicals. The mixt. of aromatic compd. and H_3PO_4 was treated, with stirring, with the alc. which was added slowly, and the mixt. was heated and stirred for 6-10 hrs.; on cooling, the org. layers were fractionated. Thus, 100 cc. benzene, 37 g. iso-BuOH, and 110 cc. H_3PO_4 (d. 1.85) gave, after 10 hrs. at 90°, 3 g. Phlu; b. 165-70°, with almost all benzene being recoverable; the use of toluene and iso-AmOH as above gave 30% mixed amyltoluenes, principally *p*-AmC₆H₄Me, b_m 204-6°. Toluene and iso-BuOH gave 56% mixed butyltoluenes, principally *p*-*tert*-BuC₆H₄Me, b_m 193-6°. Toluene and iso-PrOH gave 80% *p*-cymene; toluene and EtOH gave a very low yield of ethyltoluene, b. 157-60°. Toluene and allyl alc. gave 10% allyltoluene, b_m 183-6°, and 40% diallyltoluene, b_m 302-7°. Toluene (75 cc.), 46 g. BuCl, and 110 cc. H_3PO_4 , after 12 hrs. at 90°, gave 15% butyltoluene isomers, b. 185-95°, while *tert*-BuBr gave 37% *p*-Me₂CC₆H₄Me. PhBr (80 g.), 37 g. iso-BuOH, and 100 cc. H_3PO_4 gave, after 8 hrs. at 120-30°, 30% BuC₆H₄Br; a higher temp. gave a higher yield but also caused much tar formation; the use of PhCl gave 30% *p*-Me₂CC₆H₄Cl, b. 210-12°, while PhCl and *n*-PrOH gave 90% mixed propylchlorobenzenes, b. 185-95°, with a considerable amt. of *p*-PrC₆H₄Cl, b. 190-2°. Benzene and borneol at 70° (10 hrs.) gave 25% bornylbenzene, b_m 145-8°.

G. M. Kosolapoff

mbr., LAB. ORGANIC CHEM., MIDDLE ASIATIC STATE UNIV.,

A.S.P.-S.L.A. METALLURGICAL LITERATURE CLASSIFICATION

E.S.P. CODE CLASSIFY

E.S.P. INDEX

E.S.P. INDEX

22

10

Alkylation of aromatic compounds in the presence of phosphoric acid. II. Alkylation of phenol and anisole by acetone. V. Tambovtseva and I. Tsukryansk. *Gen. Chem. (U.S.S.R.)* 15, 830-4 (1936); cf. C.A. 40, 2707. - PhOH and PhOMe were condensed in the presence of 1.5 mols. 95% H₃PO₄ with *tert*-AmOH at 70-81°, with *iso*-PrOH and *iso*-BuOH at 80-100°, and with PhOH and BuOH at 100-30° to form 80-95% of the corresponding *p*-alkyl derivs. and sometimes a small amt. of the *o*-derivs. Below these temps., reaction did not occur. During the condensation, primary groups isomerized to secondary, and secondary to tertiary. Isolation of PhO(OBu)₂ from one reaction mixt. indicates that the reaction goes through the formation of acid esters of H₃PO₄ which are in turn the alkylating agents. With PhOMe both *Pr* and *iso*-*Pr* alic. yielded *p*-isopropylanisole, *b_m* 212-14°, *d*₄²⁰ 0.9118, *n*_D²⁰ 1.5070, and *o*-isopropylanisole, *b_m* 198-200°, *d*₄²⁰ 0.9540, *n*_D²⁰ 1.5050. Similarly, *Pr* and *iso*-*Pr* alic. and phenol yielded the corresponding *p*-*iso*-*Pr* deriv., *b_m* 213-14°, *d*₄²⁰ 1.0157, *n*_D²⁰ 1.5092, and *o*-*iso*-*Pr* deriv., crystals, *m.* 61°. Other compds. prepd. were *p*-*tert*-butylanisole, *b_m* 211-15°, *d*₄²⁰ 0.9135, *n*_D²⁰ 1.5090, and *p*-*tert*-butylphenol, crystals, *m.* 90° (from *iso*-BuOH), and *p*-*sec*-butylanisole, *b_m* 198-10°, *d*₄²⁰ 0.9110, *n*_D²⁰ 1.5050 (from BuOH). H. M. Leicester

Mem., Lab. Organic Chemistry, Middle Aortic State Univ., - 1944 -

ASAC-SLA METALLURGICAL LITERATURE CLASSIFICATION

Ed.

10

Activity of halogen in organic compounds in Friedel-Crafts syntheses. I. Condensation of ethyl esters of aliphatic halogen-substituted acids with benzene. I. P. Tsukervanik and I. V. Terent'eva. *Doklady Akad. Nauk S.S.S.R.* 50, 257-60(1945).—The position of Cl determines the nature of the products from the Friedel-Crafts reaction of C_6H_6 with esters of halo aliphatic acids; the activity of Cl rises with increased distance from the COO group. C_6H_6 (100 ml.), 40.5 g. EtO_2CCH_2Cl , and 68 g. $AlCl_3$, after standing 24 hrs., boiling 1.5 hrs., and heating 5 hrs. on a steam bath, gave 70% mixed hydrocarbons from which $EtPh$, a mixt. of isomeric $Et_2C_6H_5$, and 1,2,4- $Et_3C_6H_3$ were isolated: the latter gave 1,2,4- $C_6H_3(CO_2H)_3$, m. 215-18°, after oxidation by HNO_3 at 105-70°. C_6H_6 (100 ml.), 34 g. $EtO_2CCH_2CH_2Cl$, and 70 g. $AlCl_3$, after 24 hrs. standing and 6 hrs. at 80°, gave 68% $PhCH_2CH_2CO_2H$ (m. 49°, b. 280°), 7 g. $EtPh$, and 12 g. $PhCOCH_2CH_2Cl$ (b. 255-65°, n_D^{20} 1.4861, d_4^{20} 1.0681); merely letting the mixt. stand 14 days gave only 74% $PhCH_2CH_2CO_2H$ and $EtPh$. C_6H_6 (50 ml.), 9 g. $EtO_2C(CH_2)_2Cl$, and 20 g. $AlCl_3$, let stand 30 hrs., kept 12 hrs. at 60-70°, and boiled 3 hrs. gave 2 g. $EtPh$, 4 g. $Ph(CH_2)_2CO_2H$, m. 51°, and 4.5 g. 3,4-dihydro-1(2H)-naphthalenone, b.p. 132-4°, n_D^{20} 1.5095, d_4^{20} 1.0695. G. M. Kosolapoff

ASB-VLA METALLURGICAL LITERATURE CLASSIFICATION

FROM SYNONYMS

FROM SYNONYMS

SYNOPSIS #

SYNOPSIS #

SYNOPSIS #

SYNOPSIS #

MEL'KANOVITSKAYA, S.G.; TSUKERVANIK, I.P.

Allylation of phenol compounds in the presence of a copper catalyst. Dokl. AN Uz. SSR no. 11:40-44 '59. (MIRA 13:4)

1. Institut khimii AN UzSSR. 2. Chlen-korr. AN UzSSR (for Tsykervanik).
(Propene) (Eugenol)

SOV/7)-28-8-7/66

AUTHORS: Mel'kanovitskaya, S. G., Tsukervanik, I. P.

TITLE: Radical and Ionic Alkylation of the Aromatic Nucleus
(Radikal'noye i ionnoye alkilirovaniye aromaticheskogo yadra) VII. Butylation of Benzene, Naphthalene, Phenol, and Anisole (VII. Butilirovaniye benzola, naftalina, fenola i anizola)

PERIODICAL: Zhurnal obshchey khimii, 1958, Vol. 28, Nr 8, pp. 2032-2038 (USSR)

ABSTRACT: In connection with a previous paper (Ref 1) this paper attempts to explain the peculiarity of the alkylation of different compounds with n-butyl chloride in the presence of copper. As an example the free radical butylation of anisole was investigated. n-Butylphenyltriazeno was used as the source of free butyl radicals. In contrast to the benzylation and the reaction with diphenylchloromethane the butylation with copper proceeds much more difficultly (at 200-250°). In reacting n-butylchloride with benzene, naphthalene, phenol, and anisole the corresponding monobutyl and dibutyl aromatic compounds resulted, and in the case of phenol a butyl ether

Card 1/3

Radical and Ionic Alkylation of the Aromatic Nucleus. SOV/79-28-8-7/66
VII. Butylation of Benzene, Naphthalene, Phenol and, Anisole

of the butylphenol also resulted. Side products could not be determined. In every case it was the secondary butyl derivative which was produced. The proof of the structures was carried out by perbrominating and oxidizing. It was found that the alkylation product yield in the above reactions was 38-87 %. Thermal butylation also yielded alkyl derivatives. Under the experimental conditions used an isomerization of the primary butyl group to the secondary form always took place. The butylation of phenol follows the mechanism of indirect nuclear alkylation. Free butyl radicals produced by the thermal decomposition of n-butylphenyltriazene can be used to alkylate the anisole nucleus. There are 4 tables and 25 references, 11 of which are Soviet.

ASSOCIATION: Institut khimii Akademii nauk UzSSR (Institute of Chemistry AS UzSSR)

SUBMITTED: July 12, 1957

Card 2/3

Radical and Ionic Alkylation of the Aromatic Nucleus. RCY/79-20-8-7/66
VII. Bu~~l~~kylation of Benzene, Naphthalene, Phenol, and Anisole

Card 3/3

AUTHORS: Tsukervanik, I. P., Belinson, Z. Ya. SOV/79-28-8-8/66

TITLE: Condensations of Benzotrichloride With Benzene in the Presence of Aluminum Chloride (Kondensatsii benzotrikhlorida s benzolom v prisutstvii khloristogo alyuminiya)

PERIODICAL: Zhurnal obshchey khimii, 1958, Vol. 28, Nr 8, pp. 2038-2042 (USSR)

ABSTRACT: The authors investigated the condensations of benzotrichloride with benzene in order to produce the various substitutions of the chloride atom and to ascertain the conditions under which triphenylmethane and 9-phenylfluorene form. It was found that the amount of $AlCl_3$ present is a definite factor in this synthesis. Using 0,1 mole $AlCl_3$ (to 1 mole benzotrichloride), and independent of the temperature ($15-90^\circ$) and the reagent concentration, a yield of 70% diphenyldichloromethane was obtained with a small side product of triphenylchloromethane. With 1 mole of $AlCl_3$ triphenylchloromethane was produced as the main product with a yield of 80-98%. Further increases in the amount of $AlCl_3$ caused a hardening reaction and a decrease in the triphenylchloromethane yield. Other factors in this synthesis were studied, among them

Card 1/2